

TABLE 1A. 1998 MANAGEMENT - ON-SITE BENTGRASS TEST AT BIRMINGHAM, AL (COUNTRY CLUB OF BIRMINGHAM)

ESTABLISHMENT		FERTILIZATION			HERBICIDES		
Seeding date	16-Sep-97	Date(s)	Product	Rate (lbs./M)	Date(s)	Product	Rate (oz./M)
Problems during	None	every week, Feb. - Nov.		0.06 - 0.125 lb. of N & K20, 0.03 lb. of P205	16-Mar-98 26-May-98	Pre-San Pre-San	2 lbs. product/M 2 lbs. product/M
FACTORS OF PLAY		February	Nature Safe 8-3-5	0.5 lb. N			
		March	Nature Safe 8-3-5	0.5 lb. N			
Date opened for play		April	Nature Safe 8-3-5	0.5 lb. N			
Date closed for play	not closed	May	Nature Safe 8-3-5	0.5 lb. N			
Type of spikes allowed	softspikes	September	Nature Safe 8-3-5	0.5 lb. N			
Uses of green	chipping	October	Nature Safe 8-3-5	0.5 lb. N			
		weekly as needed	Multigreen/Maxigreen	2-4 oz./M			
MOWING		weekly as needed	Lesco Micromix & Iron	2-4 oz./M			
					INSECTICIDES		
Initial height	0.275"				Date(s)	Product	Rate (oz./M)
Current height	0.15 -0.16"						
	0.14" for short periods						
Frequency	6 days/week				as needed for cutworms	Talstar	
Type of mower	walking						
Rollers used							
Groomers used	Toro GM800 set .005" below bedknife used daily except mid-June through mid-Sept.						
		FUNGICIDES			OTHER PRODUCTS		
CULTIVATION		Date(s)	Product	Rate (oz./M)	Date(s)	Product	Rate (oz./M)
Aerification	10/13/98 using 1/2" coring times	every 2-3 wks, Feb. - mid-Dec.	Alternate Banol, Alliette,	labeled preventative rates	weekly as needed	Astron	1
Aerification	high pressure water injection every 2 wks, Feb. - Nov.		Heritage and Subdue with either Mancozeb or Daconil		weekly as needed	Perkup	3
Verticutting	lightly 2 times in spring 1998				06-May-98	Primo	8 oz./Acre
Dates of topdressing	lightly every 2 wks during growing season. Heavier after aerification.				10-Jun-98	Primo	8 oz./Acre
Other cultural practices	Rolled after coring. Rolled three times in spring 1998				NOTES/COMMENTS		

Product and company names mentioned herein may be trademarks and/or registered trademarks of their respective owners