

TABLE 11D. 2001 MANAGEMENT - ON-SITE BENTGRASS TEST AT DALLAS, TX (BENT TREE COUNTRY CLUB)

ESTABLISHMENT		FERTILIZATION			FUNGICIDES		
Seeding date	10-Sep-97	Date(s)	Product	Rate (lbs./M)	Date(s)	Product	Rate (oz./M)
Problems during	None	Granular Fertilizer Applications			Applied Heritage 2apps spring (April and May) & 2apps fall (Sept and Oct)		
FACTORS OF PLAY		February	Par Ex 16-0-24	0.5 lb. N	Application Aliette 4oz and Fore 8oz in August		
		March	Par Ex 16-0-24	0.5 lb. N	Applications of Banol @1.3oz in April/June/August and Subdue Maxx @1oz in May/July/Sept		
Date opened for play	open year round	May	Polyon 0-0-46	1 lb. K	Koban drench @ 8oz in April following Star Tine		
Date closed for play	5 days in Winter weather related plus ever Mon.	June	Polyon 0-0-46	1 lb. K	ProStar drench @6oz in May following Star Tine		
Type of spikes allowed	Softspikes	July	Polyon 0-0-46	1 lb. K			
Uses of green	Chipping and Putting	September	Polyon 0-0-46	1 lb. K			
		September	Scotts 13-2-26	0.5 lb. N	HERBICIDES		
		October	Polyon 0-0-46	1 lb. K	Date(s) Product Rate (oz./M)		
		November	Scotts 15-0-29	0.5 lb. N			
MOWING		November	Polyon 0-0-46	0.5 lb. K			
		December	Scotts 15-0-29	0.5 lb. N			
Initial height		December	Polyon 0-0-32	0.5 lb. K	Applied Conserve GC week prior to each star tine cultivation		
Current height	0.125"	Total - Granular app. 2.5 lbs. N, 0.10 lbs. P, 10.4 lbs. K					
Frequency	6 days/week						
Type of mower	Toro walkers	Foliar Fertilizer Applications			INSECTICIDES		
Rollers used	Brush 2-4X/week with solid	Every two weeks throughout year (24 Apps)			Date(s) Product Rate (oz./M)		
Groomers used	rollers all year						
CULTIVATION		Each Floratine app. consisted of:			Applied Conserve GC week prior to each star tine cultivation		
Aerfication - dates	April, June, Aug, Sept.	Total - Foliar app. 1.3 lbs. N, 0.7 lbs. P, 2.1 lbs. K			OTHER PRODUCTS		
Aerfication - type	Star tine				Date(s) Product Rate (oz./M)		
Verticutting	Verticut 1X/week - Mar-May, Oct; 1X/month - June-Sept.						
Dates of topdressing	Topdressed with sand 1X/week lightly - March-June and Sept.-Oct; sand 1X/month lightly July and August; sand medium after solid tine cultivation.	Total Fertilizer app. 3.8 lbs. N, 0.8 lbs. P, 12.5 lbs. K			None		
Other cultural practices	Triplex mount for special events (10X/year) and after Star Tine Cultivation.				NOTES/COMMENTS		

Product and company names mentioned herein may be trademarks and/or registered trademarks of their respective owners