

## **NATIONAL TURFGRASS EVALUATION PROGRAM**

The National Turfgrass Evaluation Program (NTEP) is designed to develop and coordinate uniform evaluation trials of turfgrass varieties and promising selections in the United States and Canada. Test results can be used by national companies and plant breeders to determine the broad picture of the adaptation of a cultivar. Results can also be used to determine if a cultivar is well adapted to a local area or level of turf maintenance.

Briefly, the NTEP is a self-supporting, non-profit program, sponsored by the Beltsville Agricultural Research Center and the National Turfgrass Federation, Inc. Program policy is made by a policy committee consisting of one member from each of the four (4) Regional Turfgrass Research Committees in the United States, one member from the Lawn Seed Division of the American Seed Trade Association, one member from the American Sod Producers Association, one member from the United States Golf Association (USGA) Green Section, one member from the Turfgrass Breeders Association, an executive director and a national program coordinator. The program does not make variety recommendations. However, the data from tests can be used by extension specialists and others for making recommendations.

The policy committee is responsible for determining program policy including, (1) requirements for submission of entries, (2) scheduling tests, (3) evaluation methods, (4) selecting standard or control test entries, (5) setting entry fees, (6) coordinating tests in their respective regions, (7) establishing guidelines for publication and data distribution and (8) scheduling committee meetings.

Executive Director - Kevin N. Morris, National Turfgrass Federation, Inc.

Special Projects Coordinator - Dr. Robert Shearman, University of Nebraska

### **CURRENT POLICY COMMITTEE MEMBERS:**

Mr. Craig Edminster, Cebeco International Seeds, Inc.

Dr. Thomas Fermanian, University of Illinois

Dr. Donald Floyd, Pickseed West, Inc.

Dr. Michael Kenna, USGA Green Section

Dr. Anthony Koski, Colorado State University

Dr. Peter Landschoot, Penn State University

Dr. Dennis Martin, Oklahoma State University

Dr. Jeffrey Nus, Golf Course Superintendents Assoc. of America

Mr. Ike Thomas, Turfgrass America, Inc.

### **FOR ADDITIONAL REPORTS OR INFORMATION CONTACT:**

Kevin Morris, Executive Director

National Turfgrass Evaluation Program

Beltsville Agricultural Research Center-West

Building 001, Room 245

Beltsville, Maryland 20705

[kmorris@ntep.org](mailto:kmorris@ntep.org)

[www.ntep.org](http://www.ntep.org)

CONTENTS

1996 National Buffalograss Test - 1999 data

LOCATIONS SUBMITTING DATA FOR 1999.....	1
NATIONAL BUFFALOGRASS TEST, 1996 Entries and Sponsors.....	2
Table A - 1999 Locations, Site Descriptions and Management Practices in the 1996 National Buffalograss.....	3
Table B - Locations and Data Collected in 1999.....	4
Table 1A- Mean Turfgrass Quality Ratings of Buffalograss Cultivars Grown at Eleven Locations in the U.S.....	5
Table 1B- Mean Turfgrass Quality Ratings of Buffalograss (Seeded) Cultivars Grown at Eleven Locations in the U.S.....	6
Table 1C- Mean Turfgrass Quality Ratings of Buffalograss (Vegetative) Cultivars Grown at Eleven Locations in the U.S.....	6
Table 2A- Mean Turfgrass Quality Ratings of Buffalograss Cultivars for Each Month Grown at Eleven Locations in the U.S.....	7
Table 2B- Mean Turfgrass Quality Ratings of Buffalograss (Seeded) Cultivars for Each Month Grown at Eleven Locations in the U.S.....	8
Table 2C- Mean Turfgrass Quality Ratings of Buffalograss (Vegetative) Cultivars for Each Month Grown at Eleven Locations in the U.S.....	8
Table 3A- Ranking of Mean Turfgrass Quality Ratings of Buffalograss Cultivars Grown at Eleven Locations in the U.S.....	9
Table 3B- Ranking of Mean Turfgrass Quality Ratings of Buffalograss (Seeded) Cultivars Grown at Eleven Locations in the U.S.....	10
Table 3C- Ranking of Mean Turfgrass Quality Ratings of Buffalograss (Vegetative) Cultivars Grown at Eleven Locations in the U.S.....	10
Table 4A- Genetic Color Ratings of Buffalograss Cultivars.....	11
Table 4B- Genetic Color Ratings of Buffalograss (Seeded) Cultivars.....	12
Table 4C- Genetic Color Ratings of Buffalograss (Vegetative) Cultivars.....	12
Table 5A- Spring Greenup Ratings of Buffalograss Cultivars.....	13
Table 5B- Spring Greenup Ratings of Buffalograss (Seeded) Cultivars.....	14
Table 5C- Spring Greenup Ratings of Buffalograss (Vegetative) Cultivars....	14
Table 6A- Leaf Texture Ratings of Buffalograss Cultivars.....	15

CONTENTS (continued)

Table 6B- Leaf Texture Ratings of Buffalograss (Seeded) Cultivars.....	16
Table 6C- Leaf Texture Ratings of Buffalograss (Vegetative) Cultivars.....	16
Table 7A- Spring Density Ratings of Buffalograss Cultivars.....	17
Table 7B- Spring Density Ratings of Buffalograss (Seeded) Cultivars.....	18
Table 7C- Spring Density Ratings of Buffalograss (Vegetative) Cultivars.....	18
Table 8A- Summer Density Ratings of Buffalograss Cultivars.....	19
Table 8B- Summer Density Ratings of Buffalograss (Seeded) Cultivars.....	20
Table 8C- Summer Density Ratings of Buffalograss (Vegetative) Cultivars.....	20
Table 9A- Fall Density Ratings of Buffalograss Cultivars.....	21
Table 9B- Fall Density Ratings of Buffalograss (Seeded) Cultivars.....	22
Table 9C- Fall Density Ratings of Buffalograss (Vegetative) Cultivars.....	22
Table 10A- Percent Living Ground Cover (Spring) Ratings of Buffalograss Cultivars.....	23
Table 10B- Percent Living Ground Cover (Spring) Ratings of Buffalograss (Seeded) Cultivars.....	24
Table 10C- Percent Living Ground Cover (Spring) Ratings of Buffalograss (Vegetative) Cultivars.....	24
Table 11A- Percent Living Ground Cover (Summer) Ratings of Buffalograss Cultivars.....	25
Table 11B- Percent Living Ground Cover (Summer) Ratings of Buffalograss (Seeded) Cultivars.....	26
Table 11C- Percent Living Ground Cover (Summer) Ratings of Buffalograss (Vegetative) Cultivars.....	26
Table 12A- Percent Living Ground Cover (Fall) Ratings of Buffalograss Cultivars.....	27
Table 12B- Percent Living Ground Cover (Fall) Ratings of Buffalograss (Seeded) Cultivars.....	28
Table 12C- Percent Living Ground Cover (Fall) Ratings of Buffalograss (Vegetative) Cultivars.....	28
Table 13A- Frost Tolerance Ratings of Buffalograss Cultivars.....	29
Table 13B- Frost Tolerance Ratings of Buffalograss (Seeded) Cultivars.....	30

CONTENTS (continued)

Table 13C- Frost Tolerance Ratings of Buffalograss (Vegetative) Cultivars.....	30
Table 14A- Winter Color Ratings of Buffalograss Cultivars.....	31
Table 14B- Winter Color Ratings of Buffalograss (Seeded) Cultivars.....	32
Table 14C- Winter Color Ratings of Buffalograss (Vegetative) Cultivars.....	32
Table 15A- Drought Tolerance (Dormancy) Ratings of Buffalograss Cultivars.....	33
Table 15B- Drought Tolerance (Dormancy) Ratings of Buffalograss (Seeded) Cultivars.....	34
Table 15C- Drought Tolerance (Dormancy) Ratings of Buffalograss (Vegetative) Cultivars.....	34
Table 16A- Dollar Spot Ratings of Buffalograss Cultivars.....	35
Table 16B- Dollar Spot Ratings of Buffalograss (Seeded) Cultivars.....	36
Table 16C- Dollar Spot Ratings of Buffalograss (Vegetative) Cultivars.....	36
Table 17A- Fall Color (September) Ratings of Buffalograss Cultivars.....	37
Table 17B- Fall Color (September) Ratings of Buffalograss (Seeded) Cultivars.	38
Table 17C- Fall Color (September) Ratings of Buffalograss (Vegetative) Cultivars.....	38
Table 18A- Fall Color (October) Ratings of Buffalograss Cultivars.....	39
Table 18B- Fall Color (October) Ratings of Buffalograss (Seeded) Cultivars...	40
Table 18C- Fall Color (October) Ratings of Buffalograss (Vegetative) Cultivars.....	40
Table 19A- Fall Color (November) Ratings of Buffalograss Cultivars.....	41
Table 19B- Fall Color (November) Ratings of Buffalograss (Seeded) Cultivars..	42
Table 19C- Fall Color (November) Ratings of Buffalograss (Vegetative) Cultivars.....	42
Table 20A- Fall Color (December) Ratings of Buffalograss Cultivars.....	43
Table 20B- Fall Color (December) Ratings of Buffalograss (Seeded) Cultivars..	44
Table 20C- Fall Color (December) Ratings of Buffalograss (Vegetative) Cultivars.....	44
Table 21A- Pollen Head Ratings of Buffalograss Cultivars.....	45

CONTENTS (continued)

Table 21B- Pollen Head Ratings of Buffalograss (Seeded) Cultivars.....	46
Table 21C- Pollen Head Ratings of Buffalograss (Vegetative) Cultivars.....	46
Table 22A- Percent Color (October) Ratings of Buffalograss Cultivars.....	47
Table 22B- Percent Color (October) Ratings of Buffalograss (Seeded) Cultivars.....	48
Table 22C- Percent Color (October) Ratings of Buffalograss (Vegetative) Cultivars.....	48
Table 23A- Dollar Spot Ratings of Buffalograss Cultivars at Jay, FL.....	49
Table 23B- Dollar Spot Ratings of Buffalograss (Seeded) Cultivars at Jay, FL.	50
Table 23C- Dollar Spot Ratings of Buffalograss (Vegetative) Cultivars at Jay, FL.....	50

## 1996 NATIONAL BUFFALOGRASS TEST

### LOCATIONS SUBMITTING DATA FOR 1999

<u>State</u>	<u>Location</u>	<u>Code</u>
Arizona	Tucson	AZ1
California	Riverside	CA3
Florida	Jay	FL3
Georgia	Griffin	GA1
Maryland	Silver Spring	MD1
Missouri	Columbia	MO1
Nebraska	Mead	NE1
Texas	Dallas	TX1
Texas	Lubbock	TX3
Virginia	Blacksburg	VA1
Washington	Yakima	WA4

## **1996 National Buffalograss Test**

### **Entries and Sponsors**

<b>Entry No.</b>	<b>Name</b>	<b>Type</b>	<b>Sponsor</b>
1	Cody	Seeded	Native Turf Group, Inc.
2	Tatanka	Seeded	Native Turf Group, Inc.
3	BAM-1000	Seeded	Bamert Seed Company
4	Bison	Seeded	Standard entry
5	Texoka	Seeded	Standard entry
6	91-118	Vegetative	University of Nebraska
7	86-120	Vegetative	University of Nebraska
8	Legacy (86-61)	Vegetative	Todd Valley Farms, Inc.
9	Bonnie Brae	Vegetative	Horizon Turfgrass
10	Midget	Vegetative	Horizon Turfgrass
11	Stampede	Vegetative	Turfgrass America
12	UCR-95	Vegetative	Frontier Hybrids
13	609	Vegetative	Standard entry
14	378	Vegetative	Standard entry

TABLE A.

1999 LOCATIONS, SITE DESCRIPTIONS AND MANAGEMENT PRACTICES IN  
THE 1996 NATIONAL BUFFALOGRASS TEST

LOCATION	SOIL TEXTURE	SOIL PH	SOIL PHOSPHOROUS (LBS/ACRE)	SOIL POTASSIUM (LBS/ACRE)	NITROGEN (LBS/1000 SQ FT)	SUN OR SHADE	MOWING HEIGHT (IN)	IRRIGATION PRACTICED
AZ1	SANDY LOAM	7.6-8.5	0-60	151-240	5.1-6.0	FULL SUN	2.1-2.5	TO PREVENT STRESS
CA3	SANDY LOAM	6.6-7.0	0-60	0-150	3.1-4.0	FULL SUN	1.6-2.0	TO PREVENT STRESS
FL3	SANDY LOAM	6.1-6.5	151-270	241-375	2.1-3.0	FULL SUN	1.6-2.0	TO PREVENT DORMANCY
GA1	-	-	-	-	-	-	-	-
MD1	SANDY LOAM	5.6-6.0	-	-	1.1-2.0	FULL SUN	0.0-0.5	TO PREVENT DORMANCY
MO1	SILTY CLAY LOAM	6.1-6.5	61-150	241-375	2.1-3.0	FULL SUN	1.6-2.0	TO PREVENT STRESS
NE1	-	-	-	-	-	-	-	-
TX1	SILTY CLAY AND CLAY	7.6-8.5	151-270	241-375	3.1-4.0	FULL SUN	2.1-2.5	TO PREVENT STRESS
TX3	SANDY LOAM	6.6-7.0	0-60	0-150	4.1-5.0	-	1.6-2.0	TO PREVENT STRESS
VA1	-	-	-	-	-	-	-	-
WA4	SANDY LOAM	5.6-6.0	0-60	151-240	2.1-3.0	FULL SUN	1.1-1.5	TO PREVENT STRESS

TABLE B.

## LOCATIONS AND DATA COLLECTED IN 1999

LOCATION	JANUARY QUALITY RATING	FEBRUARY QUALITY RATING	MARCH QUALITY RATING	APRIL QUALITY RATING	MAY QUALITY RATING	JUNE QUALITY RATING	JULY QUALITY RATING	AUGUST QUALITY RATING	SEPTEMBER QUALITY RATING	OCTOBER QUALITY RATING	NOVEMBER QUALITY RATING	DECEMBER QUALITY RATING	GENETIC COLOR	SPRING GREENUP	LEAF TEXTURE	SPRING DENSITY
AZ1			X	X	X	X	X	X	X	X	X	X	X			X
CA3	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
FL3			X	X	X	X	X	X	X	X	X	X	X	X	X	X
GA1				X	X	X	X	X	X	X						
MD1					X	X	X	X	X	X					X	
MO1					X	X	X	X	X	X	X	X	X	X	X	X
NE1						X	X	X	X	X	X	X	X	X	X	
TX1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
TX3			X	X	X	X	X	X	X	X	X	X	X	X	X	X
VA1				X	X	X	X	X	X	X	X		X	X	X	
WA4		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X

TABLE B. (CONT'D)

## LOCATIONS AND DATA COLLECTED IN 1999

LOCATION	SUMMER DENSITY	FALL DENSITY	PERCENT COVER SPRING	PERCENT COVER SUMMER	PERCENT COVER FALL	FROST TOLERANCE	WINTER COLOR	DROUGHT TOLERANCE	FALL DOLLAR SPOT	FALL COLOR SEPTEMBER	FALL COLOR OCTOBER	FALL COLOR NOVEMBER	FALL COLOR DECEMBER	FALL COLOR HEAD	PERCENT COLOR OCTOBER	DOLLAR SPOT APRIL	DOLLAR SPOT MAY
AZ1	X	X	X	X	X						X	X	X				
CA3							X						X	X			
FL3												X	X			X	X
GA1								X		X		X					
MD1						X											
MO1	X	X	X	X	X												
NE1										X							
TX1	X																
TX3		X					X										
VA1									X								
WA4	X	X						X				X	X				

TABLE 1A.

MEAN TURFGRASS QUALITY RATINGS OF BUFFALOGRASS CULTIVARS  
 GROWN AT ELEVEN LOCATIONS IN THE U.S. 1/  
 1999 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

NAME	AZ1	CA3	FL3	GA1	MD1	MO1	NE1	TX1	TX3	VA1	WA4	MEAN
91-118	6.8	4.9	7.7	4.2	6.4	7.2	7.2	5.7	4.9	6.2	4.1	5.9
* LEGACY (86-61)	6.6	4.3	6.2	4.2	5.4	7.0	6.3	4.1	5.0	6.5	4.9	5.5
* CODY	6.0	4.2	6.9	4.9	5.0	5.7	7.8	3.9	4.9	6.3	4.9	5.5
* TATANKA	6.4	4.4	7.0	4.6	5.1	6.2	6.7	3.4	4.8	5.8	4.9	5.4
* 609	5.9	4.6	7.4	4.7	5.4	6.5	5.6	5.6	5.3	3.8	4.3	5.4
* BONNIE BRAE	6.5	5.1	6.8	3.8	6.0	6.3	4.4	5.3	5.0	5.4	4.2	5.3
* 378	6.5	4.4	6.3	4.2	4.7	7.0	5.7	4.4	5.4	5.1	4.7	5.3
86-120	6.7	4.5	7.2	3.9	5.7	6.3	4.9	4.1	5.0	5.3	4.6	5.3
* STAMPEDE	5.8	4.5	6.8	4.8	6.2	6.8	5.1	5.4	4.7	2.9	3.9	5.2
* TEXOKA	6.5	4.1	6.4	5.0	3.8	6.4	6.4	3.9	4.8	5.2	4.0	5.1
BAM-1000	6.6	4.4	6.6	4.5	4.3	6.4	4.1	3.8	4.8	6.0	4.5	5.1
UCR-95	6.3	5.4	5.7	4.1	5.3	6.2	6.7	4.1	4.3	4.1	3.6	5.1
* BISON	6.8	4.2	6.0	4.9	3.3	6.0	6.4	4.0	4.7	4.8	3.8	5.0
* MIDGET	5.2	4.3	4.9	3.7	4.8	5.8	5.4	3.9	4.0	5.3	3.8	4.6
LSD VALUE	0.8	0.2	1.0	0.5	1.1	0.5	0.6	0.8	0.7	0.7	0.9	0.2
C.V. (%)	7.9	3.1	9.0	6.6	13.4	5.0	6.1	11.5	8.5	8.2	13.4	8.8

\* COMMERCIALLY AVAILABLE IN THE USA IN 2000.

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
 STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 1B. MEAN TURFGRASS QUALITY RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS  
GROWN AT ELEVEN LOCATIONS IN THE U.S. 1/  
1999 DATA

NAME	TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/											
	AZ1	CA3	FL3	GA1	MD1	MO1	NE1	TX1	TX3	VA1	WA4	MEAN
CODY	6.0	4.2	6.9	4.9	5.0	5.7	7.8	3.9	4.9	6.3	4.9	5.5
TATANKA	6.4	4.4	7.0	4.6	5.1	6.2	6.7	3.4	4.8	5.8	4.9	5.4
TEXOKA	6.5	4.1	6.4	5.0	3.8	6.4	6.4	3.9	4.8	5.2	4.0	5.1
BAM-1000	6.6	4.4	6.6	4.5	4.3	6.4	4.1	3.8	4.8	6.0	4.5	5.1
BISON	6.8	4.2	6.0	4.9	3.3	6.0	6.4	4.0	4.7	4.8	3.8	5.0
LSD VALUE	0.8	0.2	1.0	0.4	0.9	0.5	0.6	0.6	0.8	0.6	0.6	0.2
C.V. (%)	7.6	2.6	9.7	4.8	12.5	5.3	6.3	9.2	9.9	6.5	8.1	7.9

TABLE 1C. MEAN TURFGRASS QUALITY RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS  
GROWN AT ELEVEN LOCATIONS IN THE U.S. 1/  
1999 DATA

NAME	TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/											
	AZ1	CA3	FL3	GA1	MD1	MO1	NE1	TX1	TX3	VA1	WA4	MEAN
91-118	6.8	4.9	7.7	4.2	6.4	7.2	7.2	5.7	4.9	6.2	4.1	5.9
LEGACY (86-61)	6.6	4.3	6.2	4.2	5.4	7.0	6.3	4.1	5.0	6.5	4.9	5.5
609	5.9	4.6	7.4	4.7	5.4	6.5	5.6	5.6	5.3	3.8	4.3	5.4
BONNIE BRAE	6.5	5.1	6.8	3.8	6.0	6.3	4.4	5.3	5.0	5.4	4.2	5.3
378	6.5	4.4	6.3	4.2	4.7	7.0	5.7	4.4	5.4	5.1	4.7	5.3
86-120	6.7	4.5	7.2	3.9	5.7	6.3	4.9	4.1	5.0	5.3	4.6	5.3
STAMPEDE	5.8	4.5	6.8	4.8	6.2	6.8	5.1	5.4	4.7	2.9	3.9	5.2
UCR-95	6.3	5.4	5.7	4.1	5.3	6.2	6.7	4.1	4.3	4.1	3.6	5.1
MIDGET	5.2	4.3	4.9	3.7	4.8	5.8	5.4	3.9	4.0	5.3	3.8	4.6
LSD VALUE	0.8	0.3	0.9	0.5	1.2	0.5	0.6	0.9	0.6	0.7	1.1	0.2
C.V. (%)	8.1	3.4	8.6	7.7	13.6	4.8	6.0	12.1	7.7	9.2	15.7	9.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 2A.

MEAN TURFGRASS QUALITY RATINGS OF BUFFALOGRASS CULTIVARS FOR EACH  
MONTH GROWN AT ELEVEN LOCATIONS IN THE U.S. 1/  
1999 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF: MONTHS 2/

NAME	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	MEAN
91-118	3.8	3.0	2.8	5.5	6.2	6.9	6.9	6.6	5.8	5.6	5.0	5.2	5.9
LEGACY (86-61)	3.5	2.8	2.9	5.4	5.8	6.4	6.7	6.3	5.2	4.7	4.3	3.7	5.5
CODY	3.5	3.0	3.2	5.2	5.5	6.1	6.5	6.1	5.3	5.1	3.8	4.5	5.5
TATANKA	3.0	2.8	3.3	5.1	5.4	5.9	6.4	6.0	5.4	4.9	3.8	4.0	5.4
609	3.8	3.0	2.8	4.8	5.1	5.9	6.1	6.2	5.8	5.6	5.3	5.3	5.4
BONNIE BRAE	3.7	2.8	3.0	5.3	5.4	6.1	6.4	6.3	5.6	4.8	4.4	5.2	5.3
378	3.3	2.8	2.9	5.5	5.5	6.0	6.3	6.5	5.2	4.6	4.3	4.5	5.3
86-120	3.7	2.6	3.0	5.5	5.8	6.1	6.1	6.2	5.2	4.4	4.1	4.2	5.3
STAMPEDE	4.0	3.4	2.7	5.1	5.2	5.7	5.9	5.8	5.2	5.0	5.0	4.8	5.2
TEXOKA	3.5	2.7	3.0	5.1	5.2	5.6	6.1	5.7	5.1	5.0	3.9	4.3	5.1
EAM-1000	3.2	2.9	3.2	5.2	5.0	5.5	6.0	5.8	5.1	4.9	4.0	4.0	5.1
UCR-95	3.5	2.4	2.1	4.3	5.1	5.9	5.9	5.8	4.9	5.1	5.1	5.2	5.1
BISON	3.5	2.7	3.0	4.7	5.2	5.5	6.0	5.6	5.0	4.8	3.8	4.5	5.0
MIDGET	3.5	2.7	2.5	4.6	4.7	5.3	5.3	5.1	4.6	4.4	3.9	4.0	4.6
LSD VALUE	0.7	1.2	0.8	0.8	0.7	0.7	0.8	0.8	0.9	1.0	1.3	1.2	0.5
C.V. (%)	18.4	44.0	37.2	27.0	27.8	24.0	25.2	27.6	34.7	38.4	38.3	23.7	21.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 2B. MEAN TURFGRASS QUALITY RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS FOR EACH MONTH GROWN AT ELEVEN LOCATIONS IN THE U.S. 1/  
1999 DATA

NAME	TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF: MONTHS 2/												
	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	MEAN
CODY	3.5	3.0	3.2	5.2	5.5	6.1	6.5	6.1	5.3	5.1	3.8	4.5	5.5
TATANKA	3.0	2.8	3.3	5.1	5.4	5.9	6.4	6.0	5.4	4.9	3.8	4.0	5.4
TEXOKA	3.5	2.7	3.0	5.1	5.2	5.6	6.1	5.7	5.1	5.0	3.9	4.3	5.1
BAM-1000	3.2	2.9	3.2	5.2	5.0	5.5	6.0	5.8	5.1	4.9	4.0	4.0	5.1
BISON	3.5	2.7	3.0	4.7	5.2	5.5	6.0	5.6	5.0	4.8	3.8	4.5	5.0
LSD VALUE	0.9	1.3	0.7	0.8	0.7	0.7	0.8	0.8	0.9	1.0	1.3	0.9	0.6
C.V. (%)	25.0	50.6	30.0	26.9	28.7	26.1	25.7	29.2	34.2	38.4	41.3	17.7	21.9

TABLE 2C. MEAN TURFGRASS QUALITY RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS FOR EACH MONTH GROWN AT ELEVEN LOCATIONS IN THE U.S. 1/  
1999 DATA

NAME	TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF: MONTHS 2/												
	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	MEAN
91-118	3.8	3.0	2.8	5.5	6.2	6.9	6.9	6.6	5.8	5.6	5.0	5.2	5.9
LEGACY (86-61)	3.5	2.8	2.9	5.4	5.8	6.4	6.7	6.3	5.2	4.7	4.3	3.7	5.5
609	3.8	3.0	2.8	4.8	5.1	5.9	6.1	6.2	5.8	5.6	5.3	5.3	5.4
BONNIE BRAE	3.7	2.8	3.0	5.3	5.4	6.1	6.4	6.3	5.6	4.8	4.4	5.2	5.3
378	3.3	2.8	2.9	5.5	5.5	6.0	6.3	6.5	5.2	4.6	4.3	4.5	5.3
86-120	3.7	2.6	3.0	5.5	5.8	6.1	6.1	6.2	5.2	4.4	4.1	4.2	5.3
STAMPEDE	4.0	3.4	2.7	5.1	5.2	5.7	5.9	5.8	5.2	5.0	5.0	4.8	5.2
UCR-95	3.5	2.4	2.1	4.3	5.1	5.9	5.9	5.8	4.9	5.1	5.1	5.2	5.1
MIDGET	3.5	2.7	2.5	4.6	4.7	5.3	5.3	5.1	4.6	4.4	3.9	4.0	4.6
LSD VALUE	0.6	1.0	0.8	0.8	0.7	0.7	0.8	0.8	0.9	1.0	1.4	1.4	0.5
C.V. (%)	14.3	39.8	41.5	27.1	27.4	22.8	25.0	26.8	34.9	38.3	36.9	26.0	20.6

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 3A.

RANKING OF MEAN TURFGRASS QUALITY RATINGS OF BUFFALOGRASS CULTIVARS  
 GROWN AT ELEVEN LOCATIONS IN THE U.S. 1/  
 1999 DATA

QUALITY RANKINGS; 1=HIGHEST MEAN: STATE LOCATIONS REPORTING 2/

NAME	AZ1	CA3	FL3	GA1	MD1	MD1	NE1	TX1	TX3	VA1	WA4	MEAN
91-118	2.0	3	1	8.5	1.0	1.0	2.0	1.0	7.0	3	9	1
LEGACY (86-61)	4.5	10	11	8.5	5.5	3.0	7.0	6.5	3.0	1	3	2
CODY	11.0	13	5	3.0	9.0	14.0	1.0	12.0	6.0	2	1	3
TATANKA	9.0	7	4	6.0	8.0	10.5	3.0	14.0	10.0	5	2	4
609	12.0	4	2	5.0	5.5	5.0	9.0	2.0	2.0	13	7	5
BONNIE BRAE	7.0	2	7	13.0	3.0	8.5	13.0	4.0	4.5	6	8	6
378	8.0	8	10	10.0	11.0	2.0	8.0	5.0	1.0	10	4	7
86-120	3.0	5	3	12.0	4.0	8.5	12.0	6.5	4.5	7	5	8
STAMPEDE	13.0	6	6	4.0	2.0	4.0	11.0	3.0	11.5	14	11	9
TEXOKA	6.0	14	9	1.0	13.0	6.5	5.5	10.0	8.5	9	10	10
BAM-1000	4.5	9	8	7.0	12.0	6.5	14.0	13.0	8.5	4	6	11
UCR-95	10.0	1	13	11.0	7.0	10.5	4.0	8.0	13.0	12	14	12
BISON	1.0	12	12	2.0	14.0	12.0	5.5	9.0	11.5	11	12	13
MIDGET	14.0	11	14	14.0	10.0	13.0	10.0	11.0	14.0	8	13	14

1/ THIS TABLE CONTAINS NO STATISTICAL VALUES (LSD VALUES) THEREFORE IT SHOULD ONLY BE USED TO DETERMINE THE GENERAL PERFORMANCE OF AN ENTRY OR ENTRIES ACROSS SEVERAL LOCATIONS OR REGIONS. TO ASSESS STATISTICAL DIFFERENCES AMONG ENTRIES, REFER TO THE MEANS AND LSD VALUES FOUND IN TABLE 1.

2/ RANKING OF MEAN TURFGRASS QUALITY IS ACHIEVED BY ASSIGNING "1" TO THE HIGHEST MEAN, "2" TO THE SECOND HIGHEST MEAN, ETC. FOR EACH LOCATION. FOR EXAMPLE, IF TWO MEANS ARE TIED FOR THE SECOND AND THIRD RANKS, BOTH ARE ASSIGNED "2.5".

TABLE 3B. RANKING OF MEAN TURFGRASS QUALITY RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS  
GROWN AT ELEVEN LOCATIONS IN THE U.S. 1/  
1999 DATA

QUALITY RANKINGS; 1=HIGHEST MEAN: STATE LOCATIONS REPORTING 2/

NAME	AZ1	CA3	FL3	GA1	MD1	MO1	NE1	TX1	TX3	VA1	WA4	MEAN
CODY	5	4	2	3	2	5.0	1.0	3	1.0	1	1	1
TATANKA	4	1	1	4	1	3.0	2.0	5	4.0	3	2	2
TEXOKA	3	5	4	1	4	1.5	3.5	2	2.5	4	4	3
BAM-1000	2	2	3	5	3	1.5	5.0	4	2.5	2	3	4
BISON	1	3	5	2	5	4.0	3.5	1	5.0	5	5	5

TABLE 3C. RANKING OF MEAN TURFGRASS QUALITY RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS  
GROWN AT ELEVEN LOCATIONS IN THE U.S. 1/  
1999 DATA

QUALITY RANKINGS; 1=HIGHEST MEAN: STATE LOCATIONS REPORTING 2/

NAME	AZ1	CA3	FL3	GA1	MD1	MO1	NE1	TX1	TX3	VA1	WA4	MEAN
91-118	1	3	1	3.5	1.0	1.0	1	1.0	6.0	2	6	1
LEGACY (86-61)	3	8	7	3.5	5.5	3.0	3	6.5	3.0	1	1	2
609	7	4	2	2.0	5.5	5.0	5	2.0	2.0	8	4	3
BONNIE BRAE	4	2	5	8.0	3.0	6.5	9	4.0	4.5	3	5	4
378	5	7	6	5.0	9.0	2.0	4	5.0	1.0	6	2	5
86-120	2	5	3	7.0	4.0	6.5	8	6.5	4.5	4	3	6
STAMPEDE	8	6	4	1.0	2.0	4.0	7	3.0	7.0	9	7	7
UCR-95	6	1	8	6.0	7.0	8.0	2	8.0	8.0	7	9	8
MIDGET	9	9	9	9.0	8.0	9.0	6	9.0	9.0	5	8	9

1/ THIS TABLE CONTAINS NO STATISTICAL VALUES (LSD VALUES) THEREFORE IT SHOULD ONLY BE USED TO DETERMINE THE GENERAL PERFORMANCE OF AN ENTRY OR ENTRIES ACROSS SEVERAL LOCATIONS OR REGIONS. TO ASSESS STATISTICAL DIFFERENCES AMONG ENTRIES, REFER TO THE MEANS AND LSD VALUES FOUND IN TABLE 1.

2/ RANKING OF MEAN TURFGRASS QUALITY IS ACHIEVED BY ASSIGNING "1" TO THE HIGHEST MEAN, "2" TO THE SECOND HIGHEST MEAN, ETC. FOR EACH LOCATION. FOR EXAMPLE, IF TWO MEANS ARE TIED FOR THE SECOND AND THIRD RANKS, BOTH ARE ASSIGNED "2.5".

TABLE 4A. GENETIC COLOR RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA

NAME	GENETIC COLOR RATINGS 1-9; 9=DARK GREEN 2/								
	AZ1	CA3	FL3	MO1	NE1	TX1	VA1	WA4	MEAN
LEGACY (86-61)	8.0	7	8.0	6.0	7.0	6.0	8.0	7.0	7.1
91-118	7.7	7	8.0	6.3	8.0	5.3	6.7	5.3	6.8
BISON	8.0	7	8.0	5.3	6.7	4.0	6.3	7.3	6.6
378	8.0	7	8.0	6.0	5.7	4.7	7.3	5.7	6.5
BONNIE BRAE	7.0	7	8.0	7.0	5.0	4.3	6.0	7.0	6.4
86-120	7.7	7	8.3	5.0	6.0	5.7	6.3	5.0	6.4
CODY	7.0	7	8.0	7.0	6.0	4.7	4.0	7.0	6.3
MIDGET	6.3	7	5.7	7.0	6.3	4.7	5.3	7.0	6.2
TEXOKA	7.7	7	7.7	6.0	6.7	3.7	3.7	6.7	6.1
609	6.3	7	8.0	4.0	6.7	4.7	6.7	5.3	6.1
BAM-1000	7.7	7	7.3	6.0	7.0	3.7	3.0	7.0	6.1
TATANKA	7.3	7	7.7	6.0	5.3	4.3	3.7	7.0	6.0
STAMPEDE	6.0	7	6.3	6.7	5.0	5.3	5.7	5.7	6.0
UCR-95	6.3	6	4.0	4.0	7.0	4.3	6.0	5.3	5.4
LSD VALUE	0.8	0	1.0	0.4	0.7	1.1	1.1	1.2	0.3
C.V. (%)	7.1	0	8.1	4.5	7.3	15.2	12.6	11.5	8.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 4B. GENETIC COLOR RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA

GENETIC COLOR RATINGS 1-9; 9=DARK GREEN 2/									
NAME	AZ1	CA3	FL3	MO1	NE1	TX1	VA1	WA4	MEAN
BISON	8.0	7	8.0	5.3	6.7	4.0	6.3	7.3	6.6
CODY	7.0	7	8.0	7.0	6.0	4.7	4.0	7.0	6.3
TEXOKA	7.7	7	7.7	6.0	6.7	3.7	3.7	6.7	6.1
BAM-1000	7.7	7	7.3	6.0	7.0	3.7	3.0	7.0	6.1
TATANKA	7.3	7	7.7	6.0	5.3	4.3	3.7	7.0	6.0
LSD VALUE	1.0	0	0.7	0.4	0.7	1.5	0.7	0.6	0.3
C.V. (%)	8.4	0	5.8	4.3	7.1	22.9	10.8	5.2	8.2

TABLE 4C. GENETIC COLOR RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA

GENETIC COLOR RATINGS 1-9; 9=DARK GREEN 2/									
NAME	AZ1	CA3	FL3	MO1	NE1	TX1	VA1	WA4	MEAN
LEGACY (86-61)	8.0	7	8.0	6.0	7.0	6.0	8.0	7.0	7.1
91-118	7.7	7	8.0	6.3	8.0	5.3	6.7	5.3	6.8
378	8.0	7	8.0	6.0	5.7	4.7	7.3	5.7	6.5
BONNIE BRAE	7.0	7	8.0	7.0	5.0	4.3	6.0	7.0	6.4
86-120	7.7	7	8.3	5.0	6.0	5.7	6.3	5.0	6.4
MIDGET	6.3	7	5.7	7.0	6.3	4.7	5.3	7.0	6.2
609	6.3	7	8.0	4.0	6.7	4.7	6.7	5.3	6.1
STAMPEDE	6.0	7	6.3	6.7	5.0	5.3	5.7	5.7	6.0
UCR-95	6.3	6	4.0	4.0	7.0	4.3	6.0	5.3	5.4
LSD VALUE	0.7	0	1.1	0.4	0.8	0.9	1.3	1.4	0.3
C.V. (%)	6.1	0	9.3	4.7	7.5	10.9	12.7	14.5	9.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 5A. SPRING GREENUP RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA

NAME	SPRING GREENUP RATINGS 1-9; 9=COMPLETELY GREEN 2/								
	CA3	FL3	MD1	MO1	NE1	TX1	VA1	WA4	MEAN
378	8.0	6.0	8.0	6.0	2.7	2.7	4.3	7.0	5.6
86-120	7.3	5.7	7.7	6.0	5.3	1.0	5.3	6.0	5.5
LEGACY (86-61)	8.0	5.3	8.0	6.0	3.0	1.7	5.3	6.7	5.5
CODY	7.0	6.7	9.0	7.0	1.3	2.0	4.0	6.7	5.5
91-118	7.0	7.3	9.0	7.0	3.0	3.0	3.0	3.7	5.4
BONNIE BRAE	7.7	6.3	7.7	4.3	6.0	2.3	3.7	4.7	5.3
TATANKA	7.0	6.7	5.0	6.0	4.7	1.3	3.7	7.3	5.2
MIDGET	7.7	6.3	7.3	6.0	5.3	1.7	2.0	4.7	5.1
BISON	7.0	6.7	6.7	4.7	4.3	1.7	3.7	5.7	5.0
TEXORA	7.0	6.7	8.0	5.0	2.7	2.3	2.3	6.3	5.0
BAM-1000	7.0	7.0	7.3	6.0	1.0	2.3	3.3	5.7	5.0
STAMPEDE	7.0	7.3	7.3	4.7	2.3	1.7	1.3	5.0	4.6
609	7.0	7.0	7.3	3.0	3.0	1.3	1.0	6.3	4.5
UCR-95	6.0	7.0	6.3	3.0	2.3	1.3	1.7	4.0	4.0
LSD VALUE	0.4	1.3	2.2	1.1	2.1	1.6	1.8	2.3	0.6
C.V. (%)	3.7	12.4	18.5	12.3	39.0	52.5	35.9	25.4	21.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 5B. SPRING GREENUP RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA

NAME	SPRING GREENUP RATINGS 1-9; 9=COMPLETELY GREEN 2/								
	CA3	FL3	MD1	MO1	NE1	TX1	VA1	WA4	MEAN
CODY	7	6.7	9.0	7.0	1.3	2.0	4.0	6.7	5.5
TATANKA	7	6.7	5.0	6.0	4.7	1.3	3.7	7.3	5.2
BISON	7	6.7	6.7	4.7	4.3	1.7	3.7	5.7	5.0
TEXOKA	7	6.7	8.0	5.0	2.7	2.3	2.3	6.3	5.0
BAM-1000	7	7.0	7.3	6.0	1.0	2.3	3.3	5.7	5.0
LSD VALUE	0	1.8	3.2	0.8	1.3	1.7	2.3	0.9	0.6
C.V. (%)	0	16.7	27.5	9.0	29.2	53.4	42.3	9.1	21.3

TABLE 5C. SPRING GREENUP RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA

NAME	SPRING GREENUP RATINGS 1-9; 9=COMPLETELY GREEN 2/								
	CA3	FL3	MD1	MO1	NE1	TX1	VA1	WA4	MEAN
378	8.0	6.0	8.0	6.0	2.7	2.7	4.3	7.0	5.6
86-120	7.3	5.7	7.7	6.0	5.3	1.0	5.3	6.0	5.5
LEGACY (86-61)	8.0	5.3	8.0	6.0	3.0	1.7	5.3	6.7	5.5
91-118	7.0	7.3	9.0	7.0	3.0	3.0	3.0	3.7	5.4
BONNIE BRAE	7.7	6.3	7.7	4.3	6.0	2.3	3.7	4.7	5.3
MIDGET	7.7	6.3	7.3	6.0	5.3	1.7	2.0	4.7	5.1
STAMPEDE	7.0	7.3	7.3	4.7	2.3	1.7	1.3	5.0	4.6
609	7.0	7.0	7.3	3.0	3.0	1.3	1.0	6.3	4.5
UCR-95	6.0	7.0	6.3	3.0	2.3	1.3	1.7	4.0	4.0
LSD VALUE	0.5	0.9	1.4	1.2	2.4	1.5	1.5	2.8	0.6
C.V. (%)	4.6	8.9	11.6	14.1	41.3	52.0	30.7	32.9	20.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 6A. LEAF TEXTURE RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA

LEAF TEXTURE RATINGS 1-9; 9=VERY FINE 2/

NAME	CA3	FL3	MO1	TX1	VA1	WA4	MEAN
86-120	7.0	8.0	7.0	2.0	7.0	7.7	6.4
LEGACY (86-61)	7.7	8.0	7.0	2.0	6.3	7.7	6.4
TATANKA	7.3	7.3	7.0	2.7	7.3	7.0	6.4
BONNIE BRAE	7.0	8.0	7.7	2.0	6.3	7.0	6.3
MIDGET	7.7	7.7	5.7	2.0	7.0	8.0	6.3
UCR-95	8.0	8.0	7.0	2.0	5.3	7.3	6.3
BAM-1000	7.0	8.0	6.0	2.7	7.0	6.7	6.2
CODY	7.0	7.3	7.0	2.7	6.3	7.0	6.2
91-118	7.0	7.0	6.7	2.0	6.3	7.7	6.1
378	7.0	7.7	6.0	2.0	6.3	7.0	6.0
BISON	7.0	7.3	6.3	2.7	6.0	6.7	6.0
TEXOKA	7.0	7.7	6.0	2.3	6.0	7.0	6.0
609	7.3	7.7	6.0	3.0	4.0	7.0	5.8
STAMPEDE	7.0	7.7	7.0	2.0	3.3	7.0	5.7
LSD VALUE	0.5	0.7	0.5	0.7	1.2	0.7	0.3
C.V. (%)	4.3	5.7	4.7	19.1	12.2	6.4	7.6

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 6B. LEAF TEXTURE RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA

LEAF TEXTURE RATINGS 1-9; 9=VERY FINE 2/

NAME	CA3	FL3	MO1	TX1	VA1	WA4	MEAN
TATANKA	7.3	7.3	7.0	2.7	7.3	7.0	6.4
BAM-1000	7.0	8.0	6.0	2.7	7.0	6.7	6.2
CODY	7.0	7.3	7.0	2.7	6.3	7.0	6.2
BISON	7.0	7.3	6.3	2.7	6.0	6.7	6.0
TEXOKA	7.0	7.7	6.0	2.3	6.0	7.0	6.0
LSD VALUE	0.4	0.8	0.4	0.9	1.4	0.6	0.3
C.V. (%)	3.7	6.9	4.0	22.2	13.1	5.3	8.4

TABLE 6C. LEAF TEXTURE RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA

LEAF TEXTURE RATINGS 1-9; 9=VERY FINE 2/

NAME	CA3	FL3	MO1	TX1	VA1	WA4	MEAN
86-120	7.0	8.0	7.0	2.0	7.0	7.7	6.4
LEGACY (86-61)	7.7	8.0	7.0	2.0	6.3	7.7	6.4
BONNIE BRAE	7.0	8.0	7.7	2.0	6.3	7.0	6.3
MIDGET	7.7	7.7	5.7	2.0	7.0	8.0	6.3
UCR-95	8.0	8.0	7.0	2.0	5.3	7.3	6.3
91-118	7.0	7.0	6.7	2.0	6.3	7.7	6.1
378	7.0	7.7	6.0	2.0	6.3	7.0	6.0
609	7.3	7.7	6.0	3.0	4.0	7.0	5.8
STAMPEDE	7.0	7.7	7.0	2.0	3.3	7.0	5.7
LSD VALUE	0.5	0.6	0.5	0.5	1.1	0.8	0.3
C.V. (%)	4.6	5.0	5.0	15.8	11.5	6.9	7.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 7A. SPRING DENSITY RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	AZ1	CA3	M01	TX1	MEAN
91-118	8.0	8.7	7.3	5.0	7.3
86-120	7.0	8.3	6.3	4.7	6.6
LEGACY (86-61)	6.7	8.0	6.7	5.0	6.6
378	6.7	8.0	6.3	4.3	6.3
BONNIE BRAE	6.0	8.7	6.3	4.3	6.3
BISON	7.7	8.0	5.7	3.7	6.3
TEXOKA	7.7	7.0	6.7	3.7	6.3
CODY	6.0	7.7	7.0	4.3	6.3
609	6.3	7.7	5.7	5.0	6.2
STAMPEDE	5.7	8.0	5.7	5.3	6.2
TATANKA	6.0	8.0	6.3	4.3	6.2
BAM-1000	6.3	7.7	6.7	3.7	6.1
UCR-95	5.0	9.0	4.7	3.7	5.6
MIDGET	4.3	7.3	6.0	4.7	5.6
LSD VALUE	1.9	0.8	1.4	1.2	0.7
C.V. (%)	18.1	6.1	13.5	16.8	13.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 7B. SPRING DENSITY RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	AZ1	CA3	MO1	TX1	MEAN
BISON	7.7	8.0	5.7	3.7	6.3
TEXOKA	7.7	7.0	6.7	3.7	6.3
CODY	6.0	7.7	7.0	4.3	6.3
TATANKA	6.0	8.0	6.3	4.3	6.2
BAM-1000	6.3	7.7	6.7	3.7	6.1
LSD VALUE	1.9	0.6	1.1	1.2	0.6
C.V. (%)	17.6	4.8	10.6	18.6	12.8

TABLE 7C. SPRING DENSITY RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	AZ1	CA3	MO1	TX1	MEAN
91-118	8.0	8.7	7.3	5.0	7.3
86-120	7.0	8.3	6.3	4.7	6.6
LEGACY (86-61)	6.7	8.0	6.7	5.0	6.6
378	6.7	8.0	6.3	4.3	6.3
BONNIE BRAE	6.0	8.7	6.3	4.3	6.3
609	6.3	7.7	5.7	5.0	6.2
STAMPEDE	5.7	8.0	5.7	5.3	6.2
UCR-95	5.0	9.0	4.7	3.7	5.6
MIDGET	4.3	7.3	6.0	4.7	5.6
LSD VALUE	1.8	0.9	1.5	1.2	0.7
C.V. (%)	18.4	6.7	15.1	16.0	13.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 8A. SUMMER DENSITY RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	AZ1	MO1	TX1	WA4	MEAN
91-118	8.0	8.0	6.0	8.3	7.6
BONNIE BRAE	9.0	7.0	5.7	8.3	7.5
LEGACY (86-61)	8.3	7.7	5.0	9.0	7.5
86-120	8.0	8.0	5.0	8.7	7.4
378	8.3	7.0	4.3	9.0	7.2
609	7.0	7.7	5.3	8.3	7.1
BAM-1000	8.0	7.7	4.0	8.3	7.0
CODY	6.7	7.0	4.7	9.0	6.8
STAMPEDE	7.3	7.3	6.0	6.7	6.8
TATANKA	7.0	7.3	4.3	8.7	6.8
UCR-95	9.0	7.0	5.0	6.0	6.8
TEXOKA	7.0	7.3	4.3	8.0	6.7
BISON	7.7	7.0	4.0	7.3	6.5
MIDGET	6.0	6.3	5.0	7.7	6.3
LSD VALUE	1.5	0.7	1.0	1.7	0.6
C.V. (%)	11.9	5.6	12.2	13.2	11.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 8B. SUMMER DENSITY RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	AZ1	MO1	TX1	WA4	MEAN
BAM-1000	8.0	7.7	4.0	8.3	7.0
CODY	6.7	7.0	4.7	9.0	6.8
TATANKA	7.0	7.3	4.3	8.7	6.8
TEXOKA	7.0	7.3	4.3	8.0	6.7
BISON	7.7	7.0	4.0	7.3	6.5
LSD VALUE	1.9	0.7	1.0	0.7	0.6
C.V. (%)	15.9	6.2	14.8	5.4	10.8

TABLE 8C. SUMMER DENSITY RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	AZ1	MO1	TX1	WA4	MEAN
91-118	8.0	8.0	6.0	8.3	7.6
BONNIE BRAE	9.0	7.0	5.7	8.3	7.5
LEGACY (86-61)	8.3	7.7	5.0	9.0	7.5
86-120	8.0	8.0	5.0	8.7	7.4
378	8.3	7.0	4.3	9.0	7.2
609	7.0	7.7	5.3	8.3	7.1
STAMPEDE	7.3	7.3	6.0	6.7	6.8
UCR-95	9.0	7.0	5.0	6.0	6.8
MIDGET	6.0	6.3	5.0	7.7	6.3
LSD VALUE	1.2	0.6	0.9	2.1	0.7
C.V. (%)	9.4	5.2	11.0	16.1	11.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 9A. FALL DENSITY RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	AZ1	MO1	TX3	WA4	MEAN
609	4.0	8.7	8.3	9.0	7.5
91-118	3.3	9.0	8.7	8.7	7.4
STAMPEDE	4.0	9.0	8.3	7.3	7.2
LEGACY (86-61)	3.0	8.0	7.7	9.0	6.9
TATANKA	3.0	7.7	8.0	9.0	6.9
86-120	2.3	8.0	8.7	8.7	6.9
BONNIE BRAE	3.0	7.0	8.3	9.0	6.8
378	2.0	8.0	8.3	8.7	6.8
BAM-1000	3.7	7.0	8.0	8.3	6.8
UCR-95	5.0	8.0	7.7	6.3	6.8
CODY	2.7	7.3	7.7	8.7	6.6
BISON	3.7	7.0	7.7	7.7	6.5
TEXOKA	3.0	7.0	7.3	8.3	6.4
MIDGET	3.0	7.0	7.7	7.7	6.3
LSD VALUE	0.8	0.7	1.5	2.0	0.7
C.V. (%)	15.7	6.0	12.0	15.2	12.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 9B. FALL DENSITY RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	AZ1	MO1	TX3	WA4	MEAN
TATANKA	3.0	7.7	8.0	9.0	6.9
BAM-1000	3.7	7.0	8.0	8.3	6.8
CODY	2.7	7.3	7.7	8.7	6.6
BISON	3.7	7.0	7.7	7.7	6.5
TEXOKA	3.0	7.0	7.3	8.3	6.4
LSD VALUE	0.7	0.6	1.2	0.8	0.4
C.V. (%)	14.0	5.1	10.0	6.1	8.3

TABLE 9C. FALL DENSITY RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	AZ1	MO1	TX3	WA4	MEAN
609	4.0	8.7	8.3	9.0	7.5
91-118	3.3	9.0	8.7	8.7	7.4
STAMPEDE	4.0	9.0	8.3	7.3	7.2
LEGACY (86-61)	3.0	8.0	7.7	9.0	6.9
86-120	2.3	8.0	8.7	8.7	6.9
BONNIE BRAE	3.0	7.0	8.3	9.0	6.8
378	2.0	8.0	8.3	8.7	6.8
UCR-95	5.0	8.0	7.7	6.3	6.8
MIDGET	3.0	7.0	7.7	7.7	6.3
LSD VALUE	0.9	0.8	1.7	2.5	0.8
C.V. (%)	16.5	6.3	12.9	18.5	14.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 10A. PERCENT LIVING GROUND COVER (SPRING) RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA

PERCENT LIVING GROUND COVER IN SPRING: LOCATIONS 2/

NAME	AZ1	MOL	MEAN
91-118	80.0	83.3	81.7
TEXOKA	76.7	76.7	76.7
BISON	76.7	73.3	75.0
86-120	70.0	76.7	73.3
BAM-1000	63.3	81.7	72.5
378	66.7	76.7	71.7
LEGACY (86-61)	66.7	76.7	71.7
609	63.3	78.3	70.8
BONNIE BRAE	60.0	75.0	67.5
STAMPEDE	56.7	76.7	66.7
CODY	60.0	73.3	66.7
TATANKA	60.0	71.7	65.8
UCR-95	50.0	71.7	60.8
MIDGET	43.3	65.0	54.2
LSD VALUE	18.6	8.3	10.2
C.V. (%)	18.1	6.9	12.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 10B. PERCENT LIVING GROUND COVER (SPRING) RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA

PERCENT LIVING GROUND COVER IN SPRING: LOCATIONS 2/

NAME	AZ1	MO1	MEAN
TEXOKA	76.7	76.7	76.7
BISON	76.7	73.3	75.0
BAM-1000	63.3	81.7	72.5
CODY	60.0	73.3	66.7
TATANKA	60.0	71.7	65.8
LSD VALUE	19.0	5.9	10.0
C.V. (%)	17.6	4.8	12.3

TABLE 10C. PERCENT LIVING GROUND COVER (SPRING) RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA

PERCENT LIVING GROUND COVER IN SPRING: LOCATIONS 2/

NAME	AZ1	MO1	MEAN
91-118	80.0	83.3	81.7
86-120	70.0	76.7	73.3
378	66.7	76.7	71.7
LEGACY (86-61)	66.7	76.7	71.7
609	63.3	78.3	70.8
BONNIE BRAE	60.0	75.0	67.5
STAMPEDE	56.7	76.7	66.7
UCR-95	50.0	71.7	60.8
MIDGET	43.3	65.0	54.2
LSD VALUE	18.3	9.4	10.3
C.V. (%)	18.4	7.7	13.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 11A. PERCENT LIVING GROUND COVER (SUMMER) RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA

PERCENT LIVING GROUND COVER IN SUMMER: LOCATIONS 2/

NAME	AZ1	MOL	MEAN
BONNIE BRAE	90.0	85.0	87.5
BAM-1000	80.0	90.0	85.0
LEGACY (86-61)	83.3	85.0	84.2
UCR-95	90.0	78.3	84.2
91-118	80.0	86.7	83.3
378	83.3	81.7	82.5
86-120	80.0	83.3	81.7
BISON	76.7	85.0	80.8
STAMPEDE	73.3	85.0	79.2
TEXOKA	70.0	86.7	78.3
CODY	66.7	86.7	76.7
609	70.0	83.3	76.7
TATANKA	70.0	81.7	75.8
MIDGET	60.0	80.0	70.0
LSD VALUE	14.7	7.9	8.3
C.V. (%)	11.9	5.9	9.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 11B. PERCENT LIVING GROUND COVER (SUMMER) RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA

PERCENT LIVING GROUND COVER IN SUMMER: LOCATIONS 2/

NAME	AZ1	MO1	MEAN
BAM-1000	80.0	90.0	85.0
BISON	76.7	85.0	80.8
TEXOKA	70.0	86.7	78.3
CODY	66.7	86.7	76.7
TATANKA	70.0	81.7	75.8
LSD VALUE	18.6	6.2	9.8
C.V. (%)	15.9	4.5	10.9

TABLE 11C. PERCENT LIVING GROUND COVER (SUMMER) RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA

PERCENT LIVING GROUND COVER IN SUMMER: LOCATIONS 2/

NAME	AZ1	MO1	MEAN
BONNIE BRAE	90.0	85.0	87.5
LEGACY (86-61)	83.3	85.0	84.2
UCR-95	90.0	78.3	84.2
91-118	80.0	86.7	83.3
378	83.3	81.7	82.5
86-120	80.0	83.3	81.7
STAMPEDE	73.3	85.0	79.2
609	70.0	83.3	76.7
MIDGET	60.0	80.0	70.0
LSD VALUE	12.0	8.8	7.4
C.V. (%)	9.4	6.5	8.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 12A. PERCENT LIVING GROUND COVER (FALL) RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA

PERCENT LIVING GROUND COVER IN FALL: LOCATIONS 2/

NAME	AZ1	MD1	TX3	MEAN
609	40.0	96.3	99.0	78.4
STAMPEDE	40.0	95.7	99.0	78.2
BAM-1000	36.7	93.3	99.0	76.3
BISON	36.7	92.0	92.7	73.8
UCR-95	50.0	93.3	76.7	73.3
91-118	33.3	97.0	89.3	73.2
TEXOKA	30.0	93.7	92.7	72.1
LEGACY (86-61)	30.0	93.7	89.7	71.1
378	20.0	94.0	99.0	71.0
86-120	23.3	95.0	92.7	70.3
MIDGET	30.0	91.3	86.3	69.2
CODY	26.7	91.7	86.3	68.2
BONNIE BRAE	30.0	90.3	83.0	67.8
TATANKA	30.0	89.3	76.7	65.3
LSD VALUE	8.2	4.4	15.2	6.0
C.V. (%)	15.7	2.9	10.5	8.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 12B. PERCENT LIVING GROUND COVER (FALL) RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA

PERCENT LIVING GROUND COVER IN FALL: LOCATIONS 2/

NAME	AZ1	MO1	TX3	MEAN
BAM-1000	36.7	93.3	99.0	76.3
BISON	36.7	92.0	92.7	73.8
TEXOKA	30.0	93.7	92.7	72.1
CODY	26.7	91.7	86.3	68.2
TATANKA	30.0	89.3	76.7	65.3
LSD VALUE	7.2	5.2	14.3	5.6
C.V. (%)	14.0	3.5	9.9	8.5

TABLE 12C. PERCENT LIVING GROUND COVER (FALL) RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA

PERCENT LIVING GROUND COVER IN FALL: LOCATIONS 2/

NAME	AZ1	MO1	TX3	MEAN
609	40.0	96.3	99.0	78.4
STAMPEDE	40.0	95.7	99.0	78.2
UCR-95	50.0	93.3	76.7	73.3
91-118	33.3	97.0	89.3	73.2
LEGACY (86-61)	30.0	93.7	89.7	71.1
378	20.0	94.0	99.0	71.0
86-120	23.3	95.0	92.7	70.3
MIDGET	30.0	91.3	86.3	69.2
BONNIE BRAE	30.0	90.3	83.0	67.8
LSD VALUE	8.8	3.8	15.7	6.1
C.V. (%)	16.5	2.5	10.8	9.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 13A. FROST TOLERANCE RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA

FROST TOLERANCE RATINGS 1-9; 9=NO INJURY 2/

NAME	MD1
TATANKA	5.3
BISON	5.0
UCR-95	4.7
91-118	4.3
BONNIE BRAE	4.3
CODY	4.3
378	4.0
86-120	4.0
STAMPEDE	4.0
609	3.7
BAM-1000	3.7
LEGACY (86-61)	3.7
MIDGET	3.7
TEXOKA	3.7
LSD VALUE	2.4
C.V. (%)	36.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 13B. FROST TOLERANCE RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA

FROST TOLERANCE RATINGS 1-9; 9=NO INJURY 2/

NAME	MD1
TATANKA	5.3
BISON	5.0
CODY	4.3
BAM-1000	3.7
TEXOKA	3.7
LSD VALUE	2.7
C.V. (%)	38.5

TABLE 13C. FROST TOLERANCE RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA

FROST TOLERANCE RATINGS 1-9; 9=NO INJURY 2/

NAME	MD1
UCR-95	4.7
91-118	4.3
BONNIE BRAE	4.3
378	4.0
86-120	4.0
STAMPEDE	4.0
609	3.7
LEGACY (86-61)	3.7
MIDGET	3.7
LSD VALUE	2.3
C.V. (%)	35.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 14A. WINTER COLOR RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA

WINTER COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	CA3	WA4	MEAN
UCR-95	2	2.7	2.3
378	1	3.0	2.0
609	1	3.0	2.0
91-118	1	3.0	2.0
BAM-1000	1	3.0	2.0
BISON	1	3.0	2.0
CODY	1	3.0	2.0
LEGACY (86-61)	1	3.0	2.0
MIDGET	1	3.0	2.0
STAMPEDE	1	3.0	2.0
TATANKA	1	3.0	2.0
TEXOKA	1	3.0	2.0
86-120	1	2.3	1.7
BONNIE BRAE	1	2.3	1.7
LSD VALUE	0	0.4	0.2
C.V. (%)	0	9.3	9.6

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 14B. WINTER COLOR RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA

WINTER COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	CA3	WA4	MEAN
BAM-1000	1	3	2
BISON	1	3	2
CODY	1	3	2
TATANKA	1	3	2
TEXOKA	1	3	2
LSD VALUE	0	0	0
C.V. (%)	0	0	0

TABLE 14C. WINTER COLOR RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA

WINTER COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	CA3	WA4	MEAN
UCR-95	2	2.7	2.3
378	1	3.0	2.0
609	1	3.0	2.0
91-118	1	3.0	2.0
LEGACY (86-61)	1	3.0	2.0
MIDGET	1	3.0	2.0
STAMPEDE	1	3.0	2.0
86-120	1	2.3	1.7
BONNIE BRAE	1	2.3	1.7
LSD VALUE	0	0.5	0.3
C.V. (%)	0	11.8	12.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 15A. DROUGHT TOLERANCE (DORMANCY) RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA

DROUGHT TOLERANCE (DORMANCY) RATINGS 1-9; 9=NO DORMANCY 2/

NAME	GA1
BISON	3.0
CODY	3.0
BAM-1000	2.7
TATANKA	2.7
609	2.3
91-118	2.3
BONNIE BRAE	2.3
LEGACY (86-61)	2.3
MIDGET	2.3
UCR-95	2.3
378	2.0
86-120	2.0
STAMPEDE	2.0
TEXOKA	2.0
LSD VALUE	0.8
C.V. (%)	21.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 15B. DROUGHT TOLERANCE (DORMANCY) RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA

DROUGHT TOLERANCE (DORMANCY) RATINGS 1-9; 9=NO DORMANCY 2/

NAME	GA1
BISON	3.0
CODY	3.0
BAM-1000	2.7
TATANKA	2.7
TEXOKA	2.0
LSD VALUE	0.9
C.V. (%)	21.7

TABLE 15C. DROUGHT TOLERANCE (DORMANCY) RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA

DROUGHT TOLERANCE (DORMANCY) RATINGS 1-9; 9=NO DORMANCY 2/

NAME	GA1
609	2.3
91-118	2.3
BONNIE BRAE	2.3
LEGACY (86-61)	2.3
MIDGET	2.3
UCR-95	2.3
378	2.0
86-120	2.0
STAMPEDE	2.0
LSD VALUE	0.8
C.V. (%)	21.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 16A. DOLLAR SPOT RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA

DOLLAR SPOT RATINGS 1-9; 9=NO DISEASE 2/

NAME	GA1
609	8.0
STAMPEDE	8.0
BISON	6.5
CODY	6.0
378	5.3
UCR-95	5.3
BAM-1000	5.0
86-120	4.7
BONNIE BRAE	4.7
LEGACY (86-61)	4.7
TEXOKA	4.3
MIDGET	4.0
91-118	3.7
TATANKA	3.7
LSD VALUE	1.7
C.V. (%)	19.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 16B. DOLLAR SPOT RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA

DOLIAR SPOT RATINGS 1-9; 9=NO DISEASE 2/

NAME	GA1
BISON	6.5
CODY	6.0
BAM-1000	5.0
TEXOKA	4.3
TATANKA	3.7
LSD VALUE	2.2
C.V. (%)	26.0

TABLE 16C. DOLLAR SPOT RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA

DOLIAR SPOT RATINGS 1-9; 9=NO DISEASE 2/

NAME	GA1
609	8.0
STAMPEDE	8.0
378	5.3
UCR-95	5.3
86-120	4.7
BONNIE BRAE	4.7
LEGACY (86-61)	4.7
MIDGET	4.0
91-118	3.7
LSD VALUE	1.4
C.V. (%)	16.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 17A. FALL COLOR (SEPTEMBER) RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	AZ1	NE1	MEAN
91-118	7.3	7.3	7.3
UCR-95	6.7	7.0	6.8
BAM-1000	7.0	6.3	6.7
86-120	7.0	6.0	6.5
BISON	6.3	6.7	6.5
609	6.7	6.0	6.3
LEGACY (86-61)	6.7	6.0	6.3
MIDGET	6.0	6.7	6.3
STAMPEDE	6.7	6.0	6.3
TATANKA	7.0	5.7	6.3
TEXOKA	6.3	6.3	6.3
378	6.7	5.7	6.2
BONNIE BRAE	7.0	5.0	6.0
CODY	6.3	5.7	6.0
LSD VALUE	1.1	0.9	0.7
C.V. (%)	10.6	9.4	10.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 17B. FALL COLOR (SEPTEMBER) RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	AZ1	NE1	MEAN
BAM-1000	7.0	6.3	6.7
BISON	6.3	6.7	6.5
TATANKA	7.0	5.7	6.3
TEXOKA	6.3	6.3	6.3
CODY	6.3	5.7	6.0
LSD VALUE	1.0	0.9	0.7
C.V. (%)	9.6	9.4	9.5

TABLE 17C. FALL COLOR (SEPTEMBER) RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	AZ1	NE1	MEAN
91-118	7.3	7.3	7.3
UCR-95	6.7	7.0	6.8
86-120	7.0	6.0	6.5
609	6.7	6.0	6.3
LEGACY (86-61)	6.7	6.0	6.3
MIDGET	6.0	6.7	6.3
STAMPEDE	6.7	6.0	6.3
378	6.7	5.7	6.2
BONNIE BRAE	7.0	5.0	6.0
LSD VALUE	1.2	0.9	0.8
C.V. (%)	11.1	9.3	10.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 18A. FALL COLOR (OCTOBER) RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	AZ1	CA3	FL3	GA1	VA1	WA4	MEAN
609	5.0	6.0	7.3	6.0	7.0	5.7	6.2
STAMPEDE	5.0	6.0	6.7	4.3	7.3	5.0	5.7
UCR-95	6.0	6.0	6.0	4.3	5.7	4.7	5.4
TEXOKA	4.0	5.3	6.3	.	5.7	3.7	5.0
BISON	4.0	5.7	6.3	.	4.7	4.0	4.9
BAM-1000	4.0	5.0	6.3	.	5.0	4.0	4.9
91-118	4.3	6.0	7.7	2.7	4.3	3.0	4.7
CODY	3.7	5.0	6.3	.	4.3	3.7	4.6
MIDGET	3.7	5.3	5.7	2.3	4.0	4.7	4.3
TATANKA	4.0	5.0	5.7	.	3.0	3.7	4.3
BONNIE BRAE	3.3	7.0	6.0	2.0	3.3	2.0	3.9
86-120	2.7	7.0	6.0	2.0	3.0	2.7	3.9
LEGACY (86-61)	3.3	7.0	5.3	2.0	2.0	2.3	3.7
378	2.7	4.7	4.7	2.3	2.0	3.0	3.2
LSD VALUE	0.7	1.4	1.1	0.7	1.4	1.5	0.5
C.V. (%)	10.3	14.8	11.2	13.8	20.5	24.6	16.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 18B. FALL COLOR (OCTOBER) RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	AZ1	CA3	FL3	VA1	WA4	MEAN
TEXOKA	4.0	5.3	6.3	5.7	3.7	5.0
BISON	4.0	5.7	6.3	4.7	4.0	4.9
BAM-1000	4.0	5.0	6.3	5.0	4.0	4.9
CODY	3.7	5.0	6.3	4.3	3.7	4.6
TATANKA	4.0	5.0	5.7	3.0	3.7	4.3
LSD VALUE	0.4	1.7	1.4	1.9	1.9	0.7
C.V. (%)	6.6	20.5	13.8	26.1	31.1	20.6

TABLE 18C. FALL COLOR (OCTOBER) RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	AZ1	CA3	FL3	GA1	VA1	WA4	MEAN
609	5.0	6.0	7.3	6.0	7.0	5.7	6.2
STAMPEDE	5.0	6.0	6.7	4.3	7.3	5.0	5.7
UCR-95	6.0	6.0	6.0	4.3	5.7	4.7	5.4
91-118	4.3	6.0	7.7	2.7	4.3	3.0	4.7
MIDGET	3.7	5.3	5.7	2.3	4.0	4.7	4.3
BONNIE BRAE	3.3	7.0	6.0	2.0	3.3	2.0	3.9
86-120	2.7	7.0	6.0	2.0	3.0	2.7	3.9
LEGACY (86-61)	3.3	7.0	5.3	2.0	2.0	2.3	3.7
378	2.7	4.7	4.7	2.3	2.0	3.0	3.2
LSD VALUE	0.8	1.2	0.9	0.7	1.1	1.2	0.4
C.V. (%)	11.8	11.8	9.4	13.8	16.2	19.6	13.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 19A. FALL COLOR (NOVEMBER) RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	AZ1	CA3	FL3	VA1	MEAN
609	2.7	3.7	7.3	5.3	4.8
STAMPEDE	3.0	3.7	5.0	5.7	4.3
UCR-95	3.0	5.3	3.7	4.0	4.0
BAM-1000	2.0	2.0	3.0	3.3	2.6
91-118	2.3	1.3	4.3	2.0	2.5
TEXOKA	2.0	2.0	2.7	3.3	2.5
BISON	2.0	3.0	2.0	2.3	2.3
MIDGET	1.7	3.0	2.0	2.3	2.3
CODY	1.7	1.7	2.3	2.0	1.9
TATANKA	2.0	1.0	2.3	2.0	1.8
BONNIE BRAE	1.3	2.0	2.3	1.7	1.8
LEGACY (86-61)	1.3	1.3	1.7	2.0	1.6
86-120	1.0	1.0	1.7	1.0	1.2
378	1.0	1.0	1.3	1.0	1.1
LSD VALUE	0.6	0.9	1.4	0.7	0.5
C.V. (%)	19.6	23.4	30.2	15.0	23.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 19B. FALL COLOR (NOVEMBER) RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	AZ1	CA3	FL3	VA1	MEAN
BAM-1000	2.0	2.0	3.0	3.3	2.6
TEXOKA	2.0	2.0	2.7	3.3	2.5
BISON	2.0	3.0	2.0	2.3	2.3
CODY	1.7	1.7	2.3	2.0	1.9
TATANKA	2.0	1.0	2.3	2.0	1.8
LSD VALUE	0.4	0.8	1.6	0.7	0.5
C.V. (%)	13.4	26.7	40.5	17.2	27.7

TABLE 19C. FALL COLOR (NOVEMBER) RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	AZ1	CA3	FL3	VA1	MEAN
609	2.7	3.7	7.3	5.3	4.8
STAMPEDE	3.0	3.7	5.0	5.7	4.3
UCR-95	3.0	5.3	3.7	4.0	4.0
91-118	2.3	1.3	4.3	2.0	2.5
MIDGET	1.7	3.0	2.0	2.3	2.3
BONNIE BRAE	1.3	2.0	2.3	1.7	1.8
LEGACY (86-61)	1.3	1.3	1.7	2.0	1.6
86-120	1.0	1.0	1.7	1.0	1.2
378	1.0	1.0	1.3	1.0	1.1
LSD VALUE	0.7	0.9	1.3	0.6	0.5
C.V. (%)	22.3	21.9	25.7	13.9	22.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 20A. FALL COLOR (DECEMBER) RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	CA3
UCR-95	4.3
609	2.0
STAMPEDE	2.0
378	1.0
86-120	1.0
91-118	1.0
BAM-1000	1.0
BISON	1.0
BONNIE BRAE	1.0
CODY	1.0
LEGACY (86-61)	1.0
MIDGET	1.0
TATANKA	1.0
TEXOKA	1.0
LSD VALUE	0.2
C.V. (%)	11.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 20B. FALL COLOR (DECEMBER) RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	CA3
BAM-1000	1
BISON	1
CODY	1
TATANKA	1
TEXOKA	1
LSD VALUE	0
C.V. (%)	0

TABLE 20C. FALL COLOR (DECEMBER) RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	CA3
UCR-95	4.3
609	2.0
STAMPEDE	2.0
378	1.0
86-120	1.0
91-118	1.0
BONNIE BRAE	1.0
LEGACY (86-61)	1.0
MIDGET	1.0
LSD VALUE	0.3
C.V. (%)	12.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 21A. POLLEN HEAD RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA

POLLEN HEAD RATINGS 1-9; 9=NONE 2/

NAME	CA3
378	9.0
609	9.0
91-118	9.0
BONNIE BRAE	9.0
MIDGET	9.0
STAMPEDE	9.0
UCR-95	9.0
86-120	8.7
LEGACY (86-61)	8.3
TEXOKA	6.0
BAM-1000	5.7
BISON	5.7
CODY	5.0
TATANKA	5.0
LSD VALUE	0.7
C.V. (%)	5.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 21B. POLLEN HEAD RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA

POLLEN HEAD RATINGS 1-9; 9=NONE 2/

NAME	CA3
TEXOKA	6.0
BAM-1000	5.7
BISON	5.7
CODY	5.0
TATANKA	5.0
LSD VALUE	0.9
C.V. (%)	10.6

TABLE 21C. POLLEN HEAD RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA

POLLEN HEAD RATINGS 1-9; 9=NONE 2/

NAME	CA3
378	9.0
609	9.0
91-118	9.0
BONNIE BRAE	9.0
MIDGET	9.0
STAMPEDE	9.0
UCR-95	9.0
86-120	8.7
LEGACY (86-61)	8.3
LSD VALUE	0.4
C.V. (%)	3.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 22A. PERCENT COLOR (OCTOBER) RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA 2/

NAME	NE1
BAM-1000	8.7
CODY	6.7
91-118	4.7
609	4.0
STAMPEDE	3.7
TATANKA	3.3
MIDGET	3.0
378	2.7
86-120	2.7
BISON	2.7
BONNIE BRAE	2.7
TEXOKA	2.7
LEGACY (86-61)	2.3
UCR-95	2.0
LSD VALUE	1.1
C.V. (%)	18.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 22B. PERCENT COLOR (OCTOBER) RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA 2/

NAME	NE1
BAM-1000	8.7
CODY	6.7
TATANKA	3.3
BISON	2.7
TEXOKA	2.7
LSD VALUE	0.9
C.V. (%)	12.0

TABLE 22C. PERCENT COLOR (OCTOBER) RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA 2/

NAME	NE1
91-118	4.7
609	4.0
STAMPEDE	3.7
MIDGET	3.0
378	2.7
86-120	2.7
BONNIE BRAE	2.7
LEGACY (86-61)	2.3
UCR-95	2.0
LSD VALUE	1.2
C.V. (%)	24.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 23A. DOLLAR SPOT RATINGS OF BUFFALOGRASS CULTIVARS 1/  
1999 DATA

DOLIAR SPOT RATINGS 1-9; 9=NO DISEASE 2/

NAME	APRIL	MAY	MEAN
609	9.0	9.0	9.0
STAMPEDE	9.0	8.3	8.7
BISON	8.7	8.3	8.5
TEXOKA	8.7	8.3	8.5
UCR-95	8.7	8.3	8.5
91-118	9.0	7.3	8.2
BAM-1000	9.0	7.3	8.2
86-120	8.7	7.0	7.8
378	9.0	6.0	7.5
CODY	8.7	5.3	7.0
MIDGET	8.7	5.3	7.0
TATANKA	8.3	5.3	6.8
LEGACY (86-61)	9.0	4.0	6.5
BONNIE BRAE	5.0	4.0	4.5
LSD VALUE	1.9	4.6	2.1
C.V. (%)	11.9	31.1	14.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 23B. DOLLAR SPOT RATINGS OF BUFFALOGRASS (SEEDED) CULTIVARS 1/  
1999 DATA

DOLIAR SPOT RATINGS 1-9; 9=NO DISEASE 2/

NAME	APRIL	MAY	MEAN
BISON	8.7	8.3	8.5
TEXOKA	8.7	8.3	8.5
BAM-1000	9.0	7.3	8.2
CODY	8.7	5.3	7.0
TATANKA	8.3	5.3	6.8
LSD VALUE	-	4.2	-
C.V. (%)	5.4	27.4	13.8

TABLE 23C. DOLLAR SPOT RATINGS OF BUFFALOGRASS (VEGETATIVE) CULTIVARS 1/  
1999 DATA

DOLIAR SPOT RATINGS 1-9; 9=NO DISEASE 2/

NAME	APRIL	MAY	MEAN
609	9.0	9.0	9.0
STAMPEDE	9.0	8.3	8.7
UCR-95	8.7	8.3	8.5
91-118	9.0	7.3	8.2
86-120	8.7	7.0	7.8
378	9.0	6.0	7.5
MIDGET	8.7	5.3	7.0
LEGACY (86-61)	9.0	4.0	6.5
BONNIE BRAE	5.0	4.0	4.5
LSD VALUE	2.3	4.8	2.1
C.V. (%)	14.3	33.9	15.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.