

NATIONAL TURFGRASS EVALUATION PROGRAM

The National Turfgrass Evaluation Program (NTEP) is designed to develop and coordinate uniform evaluation trials of turfgrass varieties and promising selections in the United States and Canada. Test results can be used by national companies and plant breeders to determine the broad picture of the adaptation of a cultivar. Results can also be used to determine if a cultivar is well adapted to a local area or level of turf maintenance.

Briefly, the NTEP is a self-supporting, non-profit program, sponsored by the Beltsville Agricultural Research Center and the National Turfgrass Federation, Inc. Program policy is made by a policy committee consisting of one member from each of the four (4) Regional Turfgrass Research Committees in the United States, one member from the Lawn Seed Division of the American Seed Trade Association, one member from the United States Golf Association (USGA) Green Section, one member from the Golf Course Superintendents Assoc. of America (GCSAA), one member for the Turfgrass Producers International (TPI), one member from the Turfgrass Breeders Association and an executive director. The program does not make variety recommendations. However, the data from tests can be used by extension specialists and others for making recommendations.

The policy committee is responsible for determining program policy including, (1) requirements for submission of entries, (2) scheduling tests, (3) evaluation methods, (4) selecting standard or control test entries, (5) setting entry fees, (6) coordinating tests in their respective regions, (7) establishing guidelines for publication and data distribution and (8) scheduling committee meetings.

Executive Director - Kevin N. Morris, National Turfgrass Federation, Inc.

CURRENT POLICY COMMITTEE MEMBERS:

Mr. Craig Edminster, Cebeco International Seeds, Inc.
Dr. Donald Floyd, Pickseed West, Inc.
Dr. Michael Kenna, USGA Green Section
Dr. Peter Landschoot, Penn State University
Dr. Dennis Martin, Oklahoma State University
Dr. Gwen Stahnke, Washington State University
Mr. Ike Thomas, Turfgrass America, Inc.
Dr. Clark Throssell, Golf Course Superintendents Assoc. of America
Dr. Thomas Voigt, University of Illinois

FOR ADDITIONAL REPORTS OR INFORMATION CONTACT:

Kevin Morris, Executive Director
National Turfgrass Evaluation Program
Beltsville Agricultural Research Center-West
Building 003, Room 218
Beltsville, Maryland 20705
kmorris@ntep.org
www.ntep.org

CONTENTS

1996 National St. Augustinegrass Test - 2000 data

LOCATIONS SUBMITTING DATA FOR 2000.....1

NATIONAL ST. AUGUSTINEGRASS TEST, 1996 Entries and Sponsors.....2

Table A - 2000 Locations, Site Descriptions and Management Practices in
the 1996 National St. Augustinegrass Test.....3

Table B - Locations and Data Collected in 2000.....3

Table 1 - Mean Turfgrass Quality Ratings of St. Augustinegrass Cultivars
Grown at Six Locations in the U.S4

Table 2 - Mean Turfgrass Quality Ratings of St. Augustinegrass Cultivars
for Each Month Grown at Six Locations in the U.S.....4

Table 3 - Ranking of Mean Turfgrass Quality Ratings of St. Augustinegrass
Cultivars Grown at Six Locations in the U.S.....5

Table 4 - Genetic Color Ratings of St. Augustinegrass Cultivars.....5

Table 5 - Spring Greenup Ratings of St. Augustinegrass Cultivars.....6

Table 6 - Leaf Texture Ratings of St. Augustinegrass Cultivars.....6

Table 7 - Spring Density Ratings of St. Augustinegrass Cultivars.....7

Table 8 - Summer Density Ratings of St. Augustinegrass Cultivars.....7

Table 9 - Fall Density Ratings of St. Augustinegrass Cultivars.....8

Table 10- Percent Living Ground Cover (Spring) Ratings
of St. Augustinegrass Cultivars.....8

Table 11- Percent Living Ground Cover (Summer) Ratings
of St. Augustinegrass Cultivars.....9

Table 12- Percent Living Ground Cover (Fall) Ratings
of St. Augustinegrass Cultivars.....9

Table 13- Winter Color Ratings of St. Augustinegrass Cultivars.....10

Table 14- Percent Winter Kill Ratings of St. Augustinegrass Cultivars.....10

Table 15- Brown Patch (Warm Temperature) Ratings of
St. Augustinegrass Cultivars.....11

Table 16- Gray Leaf Spot Ratings of St. Augustinegrass Cultivars.....11

Table 17- Southern Chinch Bug Ratings of St. Augustinegrass Cultivars.....12

CONTENTS (continued)

Table 18- Fall Color (September) Ratings of St. Augustinegrass Cultivars....12

Table 19- Fall Color (October) Ratings of St. Augustinegrass Cultivars.....13

Table 20- Fall Color (November) Ratings of St. Augustinegrass Cultivars.....13

Table 21- Seedhead Ratings of St. Augustinegrass Cultivars.....14

1996 NATIONAL ST. AUGUSTINEGRASS TEST

LOCATIONS SUBMITTING DATA FOR 2000

<u>State</u>	<u>Location</u>	<u>Code</u>
Alabama	Auburn	AL1
Arizona	Tucson	AZ1
Florida	Gainesville	FL1
Louisiana	Calhoun	LA2
Mississippi	Mississippi State	MS1
North Carolina	Raleigh	NC1
Texas	College Station	TX2

1996 National St. Augustinegrass Test

Entries and Sponsors

Entry No.	Name	Sponsor
1	Delmar	Standard entry
2	Raleigh	Standard entry
3	6-89-70	The Scotts Company
4	6-89-174	The Scotts Company
5	6-89-175	The Scotts Company
6	6-89-194	The Scotts Company
7	6-89-196	The Scotts Company
8	Floritam	Standard entry

TABLE A.

2000 LOCATIONS, SITE DESCRIPTIONS AND MANAGEMENT PRACTICES IN
THE 1996 NATIONAL ST. AUGUSTINEGRASS TEST

LOCATION	SOIL TEXTURE	SOIL PH	SOIL PHOSPHOROUS (LBS/ACRE)	SOIL POTASSIUM (LBS/ACRE)	NITROGEN (LBS/1000 SQ FT)	SUN OR SHADE	MOWING HEIGHT (IN)	IRRIGATION PRACTICED
AL1	LOAMY SAND	5.6-6.0	61-150	241-375	6.1-7.0	FULL SUN	2.6-3.0	TO PREVENT STRESS
AZ1	SANDY CLAY LOAM	7.6-8.5	0-60	501+	2.1-3.0	FULL SUN	2.1-2.5	TO PREVENT STRESS
FL1	LOAMY SAND	7.1-7.5	271-450	151-240	2.1-3.0	FULL SUN	2.1-2.5	TO PREVENT STRESS
LA2	SAND	6.6-7.0	0-60	0-150	-	FULL SUN	1.1-1.5	TO PREVENT STRESS
MS1	SANDY CLAY LOAM	6.1-6.5	151-270	0-150	4.1-5.0	FULL SUN	2.1-2.5	TO PREVENT DORMANCY
NC1	SANDY CLAY LOAM	5.6-6.0	61-150	0-150	1.1-2.0	FULL SUN	2.6-3.0	TO PREVENT STRESS
TX2	SANDY LOAM	7.6-8.5	0-60	0-150	3.1-4.0	FULL SUN	2.6-3.0	TO PREVENT STRESS

TABLE B.

LOCATIONS AND DATA COLLECTED IN 2000

LOCATION	MARCH QUALITY RATING	APRIL QUALITY RATING	MAY QUALITY RATING	JUNE QUALITY RATING	JULY QUALITY RATING	AUGUST QUALITY RATING	SEPTEMBER QUALITY RATING	OCTOBER QUALITY RATING	NOVEMBER QUALITY RATING	GENETIC COLOR	SPRING GREENUP	LEAF TEXTURE	SPRING DENSITY	SUMMER DENSITY	FALL DENSITY
AL1		X	X	X	X	X	X	X		X	X	X	X	X	X
AZ1		X	X	X	X	X	X	X		X	X	X	X	X	X
FL1	X	X	X	X	X	X	X			X	X	X	X	X	X
LA2		X	X	X	X	X	X			X	X	X	X	X	X
MS1		X	X	X	X	X	X	X	X	X	X	X			
NC1			X	X	X	X	X	X	X		X				
TX2		X	X	X	X	X	X	X		X	X		X	X	X

TABLE B. (CONT'D)

LOCATIONS AND DATA COLLECTED IN 2000

LOCATION	PERCENT COVER SPRING	PERCENT COVER SUMMER	PERCENT COVER FALL	WINTER COLOR	PERCENT WINTER KILL	BROWN PATCH (WARM TEMP)	GRAY LEAF SPOT	SOUTHERN CHINCH BUG	FALL COLOR RETENTION (SEPT.)	FALL COLOR RETENTION (OCT.)	FALL COLOR RETENTION (NOV.)	SEEDHEAD RATINGS
AL1									X	X		
AZ1	X	X	X						X	X		
FL1	X	X	X	X			X					X
LA2	X	X	X									
MS1												
NC1		X						X			X	
TX2	X	X	X	X	X	X						

TABLE 1. MEAN TURFGRASS QUALITY RATINGS OF ST. AUGUSTINEGRASS CULTIVARS
GROWN AT SEVEN LOCATIONS IN THE U.S. 1/
2000 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF								
NAME	AL1	AZ1	FL1	LA2	MS1	NC1	TX2	MEAN
* DELMAR	5.0	5.8	5.2	6.5	5.7	3.9	6.0	5.5
* FLORATAM	5.4	5.4	4.1	6.8	5.4	4.9	6.0	5.4
6-89-70	5.1	4.7	5.5	.	4.8	3.6	6.2	5.0
* RALEIGH	4.5	5.0	4.0	7.3	4.6	3.9	5.6	5.0
6-89-175	3.0	6.0	3.9	.	3.1	3.5	6.0	4.2
6-89-174	4.2	5.6	1.5	.	3.0	2.3	5.4	3.7
6-89-194	3.1	5.1	1.8	.	3.2	2.5	5.2	3.5
6-89-196	2.5	4.9	2.4	.	2.8	2.8	5.2	3.4
LSD VALUE	1.4	0.8	1.7	0.6	0.9	1.4	0.8	0.5
C.V. (%)	21.5	9.2	30.4	3.9	13.8	25.4	8.3	16.7

* COMMERCIALY AVAILABLE IN THE USA IN 2001.

TABLE 2. MEAN TURFGRASS QUALITY RATINGS OF ST. AUGUSTINEGRASS CULTIVARS FOR EACH
MONTH GROWN AT SEVEN LOCATIONS IN THE U.S. 1/
2000 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF: MONTHS 2/										
NAME	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	MEAN
FLORATAM	4.0	4.9	4.9	5.2	5.3	5.5	5.8	6.0	5.7	5.4
DELMAR	5.3	5.2	5.1	5.2	5.2	5.6	5.5	5.5	5.7	5.3
6-89-70	4.0	4.3	4.1	4.8	5.2	5.6	5.5	5.5	5.7	5.0
RALEIGH	3.7	4.8	4.9	4.8	5.0	4.6	4.9	4.9	4.8	4.8
6-89-175	4.0	3.7	3.7	4.3	4.6	4.6	4.4	4.4	3.7	4.2
6-89-174	1.7	3.4	3.3	3.6	3.4	4.0	4.0	4.5	2.7	3.7
6-89-194	1.3	3.1	3.0	3.3	3.3	3.9	4.0	4.3	3.2	3.5
6-89-196	1.3	2.4	2.6	3.4	3.6	4.0	4.2	4.2	2.8	3.4
LSD VALUE	2.6	0.9	1.0	1.0	1.0	1.1	1.1	1.2	1.3	0.8
C.V. (%)	52.0	32.8	37.5	36.7	33.3	35.4	34.0	33.6	26.0	28.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 3. RANKING OF MEAN TURFGRASS QUALITY RATINGS OF ST. AUGUSTINEGRASS CULTIVARS GROWN AT SEVEN LOCATIONS IN THE U.S. 1/
2000 DATA

QUALITY RANKINGS; 1=HIGHEST MEAN; STATE LOCATIONS REPORTING 2/

NAME	AL1	AZ1	FL1	LA2	MS1	NC1	TX2	MEAN
DELMAR	3	2	2	3	1	2	2.0	1
FLORATAM	1	4	3	2	2	1	3.5	2
6-89-70	2	8	1	.	3	4	1.0	3
RALEIGH	4	6	4	1	4	3	5.0	4
6-89-175	7	1	5	.	6	5	3.5	5
6-89-174	5	3	8	.	7	8	6.0	6
6-89-194	6	5	7	.	5	7	7.5	7
6-89-196	8	7	6	.	8	6	7.5	8

1/ THIS TABLE CONTAINS NO STATISTICAL VALUES (LSD VALUES) THEREFORE IT SHOULD ONLY BE USED TO DETERMINE THE GENERAL PERFORMANCE OF AN ENTRY OR ENTRIES ACROSS SEVERAL LOCATIONS OR REGIONS. TO ASSESS STATISTICAL DIFFERENCES AMONG ENTRIES, REFER TO THE MEANS AND LSD VALUES FOUND IN TABLE 1.

2/ RANKING OF MEAN TURFGRASS QUALITY IS ACHIEVED BY ASSIGNING "1" TO THE HIGHEST MEAN, "2" TO THE SECOND HIGHEST MEAN, ETC. FOR EACH LOCATION. FOR EXAMPLE, IF TWO MEANS ARE TIED FOR THE SECOND AND THIRD RANKS, BOTH ARE ASSIGNED "2.5".

TABLE 4. GENETIC COLOR RATINGS OF ST. AUGUSTINEGRASS CULTIVARS 1/
2000 DATA

GENETIC COLOR RATINGS 1-9; 9=DARK GREEN 2/

NAME	AL1	AZ1	FL1	LA2	MS1	TX2	MEAN
DELMAR	5.0	6.0	5.7	8	7.0	7.0	6.4
6-89-196	2.7	6.3	6.3	.	8.0	8.0	6.3
6-89-175	5.3	6.0	4.7	.	6.7	8.0	6.1
6-89-70	5.7	5.0	6.0	.	7.3	6.0	6.0
6-89-174	2.7	6.7	5.7	.	7.3	7.3	5.9
RALEIGH	3.7	5.7	6.7	6	6.7	5.7	5.7
FLORATAM	3.3	6.3	6.0	6	7.0	4.7	5.6
6-89-194	2.0	6.3	5.3	.	7.0	7.0	5.5
LSD VALUE	2.3	0.7	1.4	0	0.9	1.4	0.6
C.V. (%)	38.4	7.6	15.0	0	7.6	12.9	15.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 5. SPRING GREENUP RATINGS OF ST. AUGUSTINEGRASS CULTIVARS 1/
2000 DATA

SPRING GREENUP RATINGS 1-9; 9=COMPLETELY GREEN 2/

NAME	AL1	AZ1	FL1	LA2	MS1	NC1	TX2	MEAN
DELMAR	5.7	6.0	3.3	8.0	5.7	2.0	6.0	5.2
RALEIGH	4.3	6.3	2.3	7.5	5.0	2.3	3.7	4.5
FLORATAM	5.0	5.0	3.7	6.0	4.3	1.0	4.7	4.2
6-89-175	4.0	4.0	2.7	.	4.3	2.3	4.3	3.6
6-89-70	4.0	4.0	3.0	.	4.3	1.7	3.7	3.4
6-89-174	3.7	4.0	1.3	.	4.0	2.0	4.3	3.2
6-89-196	3.3	3.0	1.3	.	3.7	2.7	3.3	2.9
6-89-194	1.7	3.3	2.3	.	4.3	1.3	2.3	2.6
LSD VALUE	2.4	2.2	1.2	4.7	0.8	1.6	1.3	0.7
C.V. (%)	37.5	30.4	29.4	28.8	11.2	51.1	20.2	28.4

TABLE 6. LEAF TEXTURE RATINGS OF ST. AUGUSTINEGRASS CULTIVARS 1/
2000 DATA

LEAF TEXTURE RATINGS 1-9; 9=VERY FINE 2/

NAME	AL1	AZ1	FL1	LA2	MS1	MEAN
6-89-175	3.0	7.3	8.0	.	7.7	6.5
6-89-196	2.0	7.7	9.0	.	7.3	6.5
6-89-174	1.3	6.7	8.3	.	6.7	5.8
6-89-194	1.0	6.7	7.3	.	7.3	5.6
6-89-70	3.0	4.7	7.0	.	7.0	5.4
DELMAR	1.7	5.7	5.7	7.0	6.3	5.3
RALEIGH	2.0	5.0	4.7	5.0	5.3	4.4
FLORATAM	1.0	4.7	2.7	4.5	4.3	3.4
LSD VALUE	1.7	0.9	1.4	3.4	1.0	0.6
C.V. (%)	55.5	8.9	12.8	27.3	9.9	15.6

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 7. SPRING DENSITY RATINGS OF ST. AUGUSTINEGRASS CULTIVARS 1/
2000 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	AL1	AZ1	FL1	LA2	TX2	MEAN
DELMAR	7.3	7.0	6.7	7.0	7.3	7.1
6-89-70	6.0	5.0	6.3	.	6.7	6.0
RALEIGH	5.7	6.3	6.7	4.0	6.7	5.9
FLORATAM	6.7	7.3	6.3	2.0	5.7	5.6
6-89-175	3.7	5.7	5.3	.	7.7	5.6
6-89-174	4.7	5.3	1.3	.	7.3	4.7
6-89-196	3.7	4.7	2.0	.	8.0	4.6
6-89-194	1.7	4.7	2.3	.	7.3	4.0
LSD VALUE	3.3	1.5	2.1	2.3	1.2	1.1
C.V. (%)	42.1	16.3	27.9	23.1	10.4	24.6

TABLE 8. SUMMER DENSITY RATINGS OF ST. AUGUSTINEGRASS CULTIVARS 1/
2000 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	AL1	AZ1	FL1	LA2	TX2	MEAN
6-89-70	8.3	4.7	6.3	.	7.7	6.8
6-89-175	2.7	7.7	4.3	.	8.0	5.7
RALEIGH	8.0	4.7	4.3	5.0	6.0	5.6
6-89-196	3.7	7.7	3.0	.	7.7	5.5
DELMAR	4.3	6.0	4.0	6.0	7.0	5.5
6-89-174	5.0	7.3	1.7	.	7.7	5.4
FLORATAM	7.7	4.7	4.0	4.5	6.0	5.4
6-89-194	1.3	7.0	2.3	.	7.7	4.6
LSD VALUE	4.4	1.1	2.3	2.5	1.0	1.2
C.V. (%)	53.0	11.4	37.3	21.6	9.0	28.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 9. FALL DENSITY RATINGS OF ST. AUGUSTINEGRASS CULTIVARS 1/
2000 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	AL1	AZ1	FL1	LA2	TX2	MEAN
6-89-70	7.0	5.7	6.3	.	7.0	6.5
DELMAR	7.0	6.7	4.0	6.0	7.3	6.2
6-89-175	2.7	8.7	4.7	.	8.0	6.0
6-89-174	5.3	7.7	2.3	.	8.0	5.8
FLORATAM	7.0	5.3	4.3	4.5	6.0	5.4
RALEIGH	3.3	5.3	4.3	7.5	5.7	5.2
6-89-196	1.0	8.3	3.0	.	8.0	5.1
6-89-194	1.7	7.3	2.3	.	8.0	4.8
LSD VALUE	3.3	0.9	2.7	3.6	0.7	1.1
C.V. (%)	46.7	8.4	42.3	26.4	6.3	24.3

TABLE 10. PERCENT LIVING GROUND COVER (SPRING) RATINGS OF ST. AUGUSTINEGRASS CULTIVARS 1/
2000 DATA

PERCENT LIVING GROUND COVER IN SPRING: LOCATIONS 2/

NAME	AZ1	FL1	LA2	TX2	MEAN
DELMAR	99.0	37.0	85.0	95.0	79.0
6-89-175	95.7	35.7	.	88.0	73.1
6-89-70	90.0	34.3	.	87.3	70.6
RALEIGH	99.0	29.0	40.0	89.0	64.3
6-89-174	94.7	12.3	.	84.3	63.8
FLORATAM	99.0	45.7	20.0	86.7	62.8
6-89-196	70.0	14.0	.	93.3	59.1
6-89-194	81.0	23.0	.	68.3	57.4
LSD VALUE	18.4	18.0	22.7	13.2	9.3
C.V. (%)	12.5	38.8	20.7	9.5	15.6

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 11. PERCENT LIVING GROUND COVER (SUMMER) RATINGS OF ST. AUGUSTINEGRASS CULTIVARS 1/
2000 DATA

PERCENT LIVING GROUND COVER IN SUMMER: LOCATIONS 2/

NAME	AZ1	FL1	LA2	NC1	TX2	MEAN
DELMAR	95.3	60.0	90.0	88.3	98.3	86.4
RALEIGH	95.3	62.3	65.0	96.0	97.7	83.3
FLORATAM	92.3	68.7	32.5	97.7	97.7	77.8
6-89-70	80.0	48.7	.	82.3	97.0	77.0
6-89-175	90.0	46.3	.	70.0	93.0	74.8
6-89-174	80.0	10.0	.	38.3	97.0	56.3
6-89-196	53.3	24.0	.	57.7	89.3	56.1
6-89-194	70.0	25.3	.	21.7	94.3	52.8
LSD VALUE	21.6	26.7	57.2	43.4	11.3	14.0
C.V. (%)	16.4	38.5	40.2	39.1	7.3	24.9

TABLE 12. PERCENT LIVING GROUND COVER (FALL) RATINGS OF ST. AUGUSTINEGRASS CULTIVARS 1/
2000 DATA

PERCENT LIVING GROUND COVER IN FALL: LOCATIONS 2/

NAME	AZ1	FL1	LA2	TX2	MEAN
DELMAR	97.7	71.7	90.0	97.7	89.3
RALEIGH	99.0	79.0	65.0	96.3	84.8
6-89-70	92.3	55.7	.	97.0	81.7
6-89-175	96.3	51.7	.	95.7	81.2
FLORATAM	97.7	80.3	32.5	99.0	77.4
6-89-194	89.7	26.3	.	97.0	71.0
6-89-196	81.7	28.7	.	95.7	68.7
6-89-174	89.7	9.0	.	97.0	65.2
LSD VALUE	8.3	35.4	57.2	2.9	12.4
C.V. (%)	5.5	43.8	40.2	1.8	17.6

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 13. WINTER COLOR RATINGS OF ST. AUGUSTINEGRASS CULTIVARS 1/
2000 DATA

WINTER COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	FL1	TX2	MEAN
DELMAR	4.7	7.7	6.2
FLORATAM	4.0	8.0	6.0
RALEIGH	3.7	7.3	5.5
6-89-70	3.3	7.0	5.2
6-89-175	2.7	7.3	5.0
6-89-194	1.3	6.3	3.8
6-89-174	1.3	5.0	3.2
6-89-196	1.0	5.0	3.0
LSD VALUE	1.3	0.9	0.8
C.V. (%)	28.7	8.1	14.3

TABLE 14. PERCENT WINTER KILL RATINGS OF ST. AUGUSTINEGRASS CULTIVARS 1/
2000 DATA

PERCENT WINTER KILL RATINGS: LOCATIONS 2/

NAME	TX2
6-89-194	31.7
6-89-174	15.7
FLORATAM	13.3
6-89-70	12.7
6-89-175	12.0
RALEIGH	11.0
6-89-196	6.7
DELMAR	5.0
LSD VALUE	13.2
C.V. (%)	61.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 15. BROWN PATCH (WARM TEMPERATURE) RATINGS OF ST. AUGUSTINEGRASS CULTIVARS 1/
2000 DATA

BROWN PATCH RATINGS 1-9; 9=NO DISEASE 2/

NAME	TX2
FLORATAM	7.7
6-89-194	7.3
6-89-196	7.3
6-89-174	7.0
6-89-175	6.7
6-89-70	5.7
DELMAR	5.3
RALEIGH	5.3
LSD VALUE	2.5
C.V. (%)	23.8

TABLE 16. GRAY LEAF SPOT RATINGS OF ST. AUGUSTINEGRASS CULTIVARS 1/
2000 DATA

GRAY LEAF SPOT RATINGS 1-9; 9=NO DISEASE 2/

NAME	FL1
6-89-70	9.0
RALEIGH	9.0
6-89-175	8.3
DELMAR	8.3
6-89-194	7.7
6-89-196	7.7
FLORATAM	6.7
6-89-174	6.0
LSD VALUE	1.9
C.V. (%)	14.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 17. SOUTHERN CHINCH BUG RATINGS OF ST. AUGUSTINEGRASS CULTIVARS 1/
2000 DATA

SOUTHERN CHINCH BUG RATINGS 1-9; 9=NO DAMAGE 2/

NAME	NC1
FLORATAM	9.0
6-89-70	8.3
6-89-196	7.3
RALEIGH	7.3
6-89-174	7.0
6-89-194	6.3
DELMAR	6.3
6-89-175	6.0
LSD VALUE	1.9
C.V. (%)	16.0

TABLE 18. FALL COLOR (SEPTEMBER) RATINGS OF ST. AUGUSTINEGRASS CULTIVARS 1/
2000 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	AL1	AZ1	MEAN
6-89-175	5.0	6.7	5.8
DELMAR	5.7	5.7	5.7
6-89-196	3.3	7.7	5.5
6-89-174	4.0	6.7	5.3
FLORATAM	4.0	6.3	5.2
6-89-70	4.3	5.7	5.0
6-89-194	2.3	7.0	4.7
RALEIGH	4.3	4.3	4.3
LSD VALUE	3.0	1.0	1.6
C.V. (%)	45.9	10.3	27.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 19. FALL COLOR (OCTOBER) RATINGS OF ST. AUGUSTINEGRASS CULTIVARS 1/
2000 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	AL1	AZ1	MEAN
6-89-174	6.0	7.7	6.8
DELMAR	5.7	6.7	6.2
6-89-70	4.3	6.3	5.3
RALEIGH	5.3	5.3	5.3
FLORATAM	3.7	6.7	5.2
6-89-194	3.0	7.3	5.2
6-89-175	2.7	7.3	5.0
6-89-196	2.0	8.0	5.0
LSD VALUE	3.0	1.6	1.7
C.V. (%)	45.5	14.8	27.3

TABLE 20. FALL COLOR (NOVEMBER) RATINGS OF ST. AUGUSTINEGRASS CULTIVARS 1/
2000 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	NC1
6-89-70	5.0
FLORATAM	5.0
6-89-174	4.7
6-89-194	4.3
6-89-196	4.3
RALEIGH	4.3
6-89-175	4.0
DELMAR	4.0
LSD VALUE	0.7
C.V. (%)	9.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 21. SEEDHEAD RATINGS OF ST. AUGUSTINEGRASS CULTIVARS 1/
2000 DATA

SEEDHEAD RATINGS 1-9; 9=NONE 2/

NAME	FL1
6-89-196	8.7
6-89-194	7.7
FLORATAM	7.7
DELMAR	6.0
6-89-174	5.3
6-89-70	4.3
6-89-175	4.0
RALEIGH	1.0
LSD VALUE	2.5
C.V. (%)	27.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.