

## **NATIONAL TURFGRASS EVALUATION PROGRAM**

The National Turfgrass Evaluation Program (NTEP) is designed to develop and coordinate uniform evaluation trials of turfgrass varieties and promising selections in the United States and Canada. Test results can be used by national companies and plant breeders to determine the broad picture of the adaptation of a cultivar. Results can also be used to determine if a cultivar is well adapted to a local area or level of turf maintenance.

Briefly, the NTEP is a self-supporting, non-profit program, sponsored by the Beltsville Agricultural Research Center and the National Turfgrass Federation, Inc. Program policy is made by a policy committee consisting of one member from each of the four (4) Regional Turfgrass Research Committees in the United States, one member from the Lawn Seed Division of the American Seed Trade Association, one member from the United States Golf Association (USGA) Green Section, one member from the Golf Course Superintendents Assoc. of America (GCSAA), one member for the Turfgrass Producers International (TPI), one member from the Turfgrass Breeders Association and an executive director. The program does not make variety recommendations. However, the data from tests can be used by extension specialists and others for making recommendations.

The policy committee is responsible for determining program policy including, (1) requirements for submission of entries, (2) scheduling tests, (3) evaluation methods, (4) selecting standard or control test entries, (5) setting entry fees, (6) coordinating tests in their respective regions, (7) establishing guidelines for publication and data distribution and (8) scheduling committee meetings.

Executive Director - Kevin N. Morris, National Turfgrass Federation, Inc.

### **CURRENT POLICY COMMITTEE MEMBERS:**

Dr. Steve Johnson, DLF International Seeds  
Ms. Chris McDowell, Pickseed West, Inc.  
Dr. Michael Kenna, USGA Green Section  
Dr. David Williams, University of Kentucky  
Dr. Bernd Leinauer, New Mexico State University  
Mr. Ike Thomas  
Dr. Clark Throssell, Golf Course Superintendents Assoc. of America  
Dr. Brian Horgan, University of Minnesota  
Mr. Duane Klundt, Turf-Seed, Inc.

### **FOR ADDITIONAL REPORTS OR INFORMATION CONTACT:**

Kevin Morris, Executive Director  
National Turfgrass Evaluation Program  
Beltsville Agricultural Research Center-West  
Building 003, Room 218  
Beltsville, Maryland 20705  
[kmorris@ntep.org](mailto:kmorris@ntep.org)  
[www.ntep.org](http://www.ntep.org)

CONTENTS

2002 National Zoysiagrass Test - 2005 data

LOCATIONS SUBMITTING DATA FOR 2005.....	1
NATIONAL ZOYSIAGRASS TEST, 2002 Entries and Sponsors.....	2
Table A - 2005 Locations, Site Descriptions and Management Practices in the 2002 National Zoysiagrass Test.....	3
Table B - Locations and Data Collected in 2005.....	4
Table 1A - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars Grown at Eleven Locations in the U.S. Maintained using "Schedule A".....	7
Table 1B - Mean Turfgrass Quality Ratings of Zoysiagrass (Seeded) Cultivars Grown at Eleven Locations in the U.S. Maintained using "Schedule A".....	8
Table 1C - Mean Turfgrass Quality Ratings of Zoysiagrass (Vegetative) Cultivars Grown at Eleven Locations in the U.S. Maintained using "Schedule A".....	8
Table 2A - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars Grown at Five Locations in the U.S. Maintained using "Schedule B".....	9
Table 2B - Mean Turfgrass Quality Ratings of Zoysiagrass (Seeded) Cultivars Grown at Five Locations in the U.S. Maintained using "Schedule B".....	10
Table 2C - Mean Turfgrass Quality Ratings of Zoysiagrass (Vegetative) Cultivars Grown at Five Locations in the U.S. Maintained using "Schedule B".....	10
Table 3A - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars Grown at Eight Locations in the Transition Region.....	11
Table 3B - Mean Turfgrass Quality Ratings of Zoysiagrass (Seeded) Cultivars Grown at Eight Locations in the Transition Region.....	12
Table 3C - Mean Turfgrass Quality Ratings of Zoysiagrass (Vegetative) Cultivars Grown at Eight Locations in the Transition Region.....	12
Table 4A - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars Grown at Five Locations in the Southeast Region.....	13
Table 4B - Mean Turfgrass Quality Ratings of Zoysiagrass (Seeded) Cultivars Grown at Five Locations in the Southeast Region.....	14
Table 4C - Mean Turfgrass Quality Ratings of Zoysiagrass (Vegetative) Cultivars Grown at Five Locations in the Southeast Region.....	14

CONTENTS (continued)

Table 5A - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars for Each Month Grown at West Lafayette, IN.....	15
Table 5B - Mean Turfgrass Quality Ratings of Zoysiagrass (Seeded) Cultivars for Each Month Grown at West Lafayette, IN.....	16
Table 5C - Mean Turfgrass Quality Ratings of Zoysiagrass (Vegetative) Cultivars for Each Month Grown at West Lafayette, IN.....	16
Table 6A - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars Grown at Two Locations in the Southwest Region.....	17
Table 6B - Mean Turfgrass Quality Ratings of Zoysiagrass (Seeded) Cultivars Grown at Two Locations in the Southwest Region.....	18
Table 6C - Mean Turfgrass Quality Ratings of Zoysiagrass (Vegetative) Cultivars Grown at Two Locations in the Southwest Region.....	18
Table 7A - Genetic Color Ratings of Zoysiagrass Cultivars.....	19
Table 7B - Genetic Color Ratings of Zoysiagrass (Seeded) Cultivars.....	20
Table 7C - Genetic Color Ratings of Zoysiagrass (Vegetative) Cultivars.....	20
Table 8A - Spring Greenup Ratings of Zoysiagrass Cultivars.....	21
Table 8B - Spring Greenup Ratings of Zoysiagrass (Seeded) Cultivars.....	22
Table 8C - Spring Greenup Ratings of Zoysiagrass (Vegetative) Cultivars.....	22
Table 9A - Leaf Texture Ratings of Zoysiagrass Cultivars.....	23
Table 9B - Leaf Texture Ratings of Zoysiagrass (Seeded) Cultivars.....	24
Table 9C - Leaf Texture Ratings of Zoysiagrass (Vegetative) Cultivars.....	24
Table 10A- Spring Density Ratings of Zoysiagrass Cultivars.....	25
Table 10B- Spring Density Ratings of Zoysiagrass (Seeded) Cultivars.....	26
Table 10C- Spring Density Ratings of Zoysiagrass (Vegetative) Cultivars.....	26
Table 11A- Summer Density Ratings of Zoysiagrass Cultivars.....	27
Table 11B- Summer Density Ratings of Zoysiagrass (Seeded) Cultivars.....	28
Table 11C- Summer Density Ratings of Zoysiagrass (Vegetative) Cultivars.....	28
Table 12A- Fall Density Ratings of Zoysiagrass Cultivars.....	29
Table 12B- Fall Density Ratings of Zoysiagrass (Seeded) Cultivars.....	30
Table 12C- Fall Density Ratings of Zoysiagrass (Vegetative) Cultivars.....	30

CONTENTS (continued)

Table 13A- Percent Living Ground Cover (Spring) Ratings of Zoysiagrass Cultivars.....	31
Table 13B- Percent Living Ground Cover (Spring) Ratings of Zoysiagrass (Seeded) Cultivars.....	32
Table 13C- Percent Living Ground Cover (Spring) Ratings of Zoysiagrass (Vegetative) Cultivars.....	32
Table 14A- Percent Living Ground Cover (Summer) Ratings of Zoysiagrass Cultivars.....	33
Table 14B- Percent Living Ground Cover (Summer) Ratings of Zoysiagrass (Seeded) Cultivars.....	34
Table 14C- Percent Living Ground Cover (Summer) Ratings of Zoysiagrass (Vegetative) Cultivars.....	34
Table 15A- Percent Living Ground Cover (Fall) Ratings of Zoysiagrass Cultivars.....	35
Table 15B- Percent Living Ground Cover (Fall) Ratings of Zoysiagrass (Seeded) Cultivars.....	36
Table 15C- Percent Living Ground Cover (Fall) Ratings of Zoysiagrass (Vegetative) Cultivars.....	36
Table 16A- Frost Tolerance Ratings of Zoysiagrass Cultivars.....	37
Table 16B- Frost Tolerance Ratings of Zoysiagrass (Seeded) Cultivars.....	38
Table 16C- Frost Tolerance Ratings of Zoysiagrass (Vegetative) Cultivars.....	38
Table 17A- Winter Color Ratings of Zoysiagrass Cultivars.....	39
Table 17B- Winter Color Ratings of Zoysiagrass (Seeded) Cultivars.....	40
Table 17C- Winter Color Ratings of Zoysiagrass (Vegetative) Cultivars.....	40
Table 18A- Dollar Spot Ratings of Zoysiagrass Cultivars.....	41
Table 18B- Dollar Spot Ratings of Zoysiagrass (Seeded) Cultivars.....	42
Table 18C- Dollar Spot Ratings of Zoysiagrass (Vegetative) Cultivars.....	42
Table 19A- Brown Patch (Warm Temperature) Ratings of Zoysiagrass Cultivars.....	43
Table 19B- Brown Patch (Warm Temperature) Ratings of Zoysiagrass (Seeded) Cultivars.....	44
Table 19C- Brown Patch (Warm Temperature) Ratings of Zoysiagrass (Vegetative) Cultivars.....	44
Table 20A- Take-All Patch Ratings of Zoysiagrass Cultivars.....	45

CONTENTS (continued)

Table 20B- Take-All Patch Ratings of Zoysiagrass (Seeded) Cultivars.....	46
Table 20C- Take-All Patch Ratings of Zoysiagrass (Vegetative) Cultivars.....	46
Table 21A- Mole Cricket Ratings of Zoysiagrass Cultivars.....	47
Table 21B- Mole Cricket Ratings of Zoysiagrass (Seeded) Cultivars.....	48
Table 21C- Mole Cricket Ratings of Zoysiagrass (Vegetative) Cultivars.....	48
Table 22A- Fall Color (September) Ratings of Zoysiagrass Cultivars.....	49
Table 22B- Fall Color (September) Ratings of Zoysiagrass (Seeded) Cultivars....	50
Table 22C- Fall Color (September) Ratings of Zoysiagrass (Vegetative) Cultivars.....	50
Table 23A- Fall Color (October) Ratings of Zoysiagrass Cultivars.....	51
Table 23B- Fall Color (October) Ratings of Zoysiagrass (Seeded) Cultivars.....	52
Table 23C- Fall Color (October) Ratings of Zoysiagrass (Vegetative) Cultivars.....	52
Table 24A- Fall Color (November) Ratings of Zoysiagrass Cultivars.....	53
Table 24B- Fall Color (November) Ratings of Zoysiagrass (Seeded) Cultivars....	54
Table 24C- Fall Color (November) Ratings of Zoysiagrass (Vegetative) Cultivars.....	54
Table 25A- Fall Color (December) Ratings of Zoysiagrass Cultivars.....	55
Table 25B- Fall Color (December) Ratings of Zoysiagrass (Seeded) Cultivars....	56
Table 25C- Fall Color (December) Ratings of Zoysiagrass (Vegetative) Cultivars.....	56
Table 26A- Seedhead Ratings of Zoysiagrass Cultivars.....	57
Table 26B- Seedhead Ratings of Zoysiagrass (Seeded) Cultivars.....	58
Table 26C- Seedhead Ratings of Zoysiagrass (Vegetative) Cultivars.....	58
Table 27A- Mowing Quality Ratings of Zoysiagrass Cultivars.....	59
Table 27B- Mowing Quality Ratings of Zoysiagrass (Seeded) Cultivars.....	60
Table 27C- Mowing Quality Ratings of Zoysiagrass (Vegetative) Cultivars.....	60
Table 28A- Mean Large Zoysia Patch Measurements of Zoysiagrass Cultivars .....	61
Table 28B- Mean Large Zoysia Patch Measurements of Zoysiagrass (Seeded) Cultivars.....	62

**CONTENTS (continued)**

<b>Table 28C- Mean Large Zoysia Patch Measurements of Zoysiagrass (Vegetative) Cultivars.....</b>	<b>62</b>
<b>Table 29A- Large Patch Ratings of Zoysiagrass Cultivars at Raleigh, NC.....</b>	<b>63</b>
<b>Table 29B- Large Patch Ratings of Zoysiagrass (Seeded) Cultivars at Raleigh, NC.....</b>	<b>64</b>
<b>Table 29C- Large Patch Ratings of Zoysiagrass (Vegetative) Cultivars at Raleigh, NC.....</b>	<b>64</b>
<b>Appendix Table A - Summary of Turfgrass Quality Ratings of Zoysiagrass Cultivars.....</b>	<b>65</b>

## A Guide to NTEP Turfgrass Ratings

### Introduction

The quality and scientific merit of NTEP data is extremely important. However, the evaluation of turfgrass species and cultivars is a difficult and complex issue. Furthermore, turfgrass evaluation is generally a subjective process based on visual estimates of factors, like genetic color, stand density, leaf texture, uniformity and quality. These factors can not be measured in the same way as other agricultural crops. Turfgrass quality is not a measure of yield or nutritive value. Turfgrass quality is a measure of aesthetics (i.e. density, uniformity, texture, smoothness, growth habit and color), and functional use. The most common way of assessing turfgrass quality is a visual rating system that is based on the turfgrass evaluator's judgement.

### General Considerations

Most visual ratings collected on NTEP trials are based on a 1 to 9 rating scale. One is the poorest or lowest and 9 is the best or highest rating. However, a few characteristics, such as winter kill or percent living ground cover, are rated on a percentage basis, again by using the evaluator's judgement. Most disease ratings found in NTEP reports will use the 1-9 scale, 9=no disease except where the evaluator made a judgement of the percentage of disease in each plot. Percent disease data will be found in separate tables and will normally not be included with disease data using the 1-9 scale.

### Turfgrass Quality

Turfgrass Quality is based on 9 being outstanding or ideal turf and 1 being poorest or dead. A rating of 6 or above is generally considered acceptable. A quality rating value of 9 is reserved for a perfect or ideal grass, but it also can reflect an absolutely outstanding treatment plot. The NTEP requires quality ratings on a monthly basis. Quality ratings take into account the aesthetic and functional aspects of the turf. Quality ratings are not based on color alone, but on a combination of color, density, uniformity, texture, and disease or environmental stress.

Turfgrass quality ratings are grouped and presented by region, management level, a particular stress (shade, traffic, etc.) and in some cases, by individual location (starting with 2001 data, data from each location will be posted separately as well on the NTEP web site, <http://www.ntep.org>). Also available now is a summary table (Appendix) in the back of this report. This summary table includes various statistical measures not previously compiled for NTEP reports. For an explanation of this table and these changes, please go to the NTEP web site at <http://www.ntep.org/pdf/grandmean.mem.pdf>.

### Other Ratings

More detailed information on the ratings of specific characteristics can be found on the NTEP web site at <http://www.ntep.org/reports/ratings.htm>.

**2002 NATIONAL ZOYSIAGRASS TEST**

**LOCATIONS SUBMITTING DATA FOR 2005**

<u>State</u>	<u>Location</u>	<u>Code</u>
Arkansas	Fayetteville	AR1
California	Riverside	CA3
Florida	Jay	FL3
Georgia	Griffin	GA1
Illinois	Carbondale	IL2
Indiana	West Lafayette	IN1
Kansas	Manhattan	KS1
Kentucky	Lexington	KY1
Maryland	College Park	MD1
Mississippi	Mississippi State	MS1
Missouri	Columbia	MO1
New Mexico	Las Cruces	NM1
North Carolina	Raleigh	NC1
Oklahoma	Stillwater	OK1
South Carolina	Florence	SC1
Texas	Dallas	TX1

## 2002 National Zoysiagrass Test

### Entries and Sponsors

Entry No.	Name	Type	Sponsor
1	Meyer	Vegetative	Standard entry
2	Emerald	Vegetative	Standard entry
3	Himeno	Vegetative	Zoysian Japan Co.
4	J-37	Seeded	Standard entry
5	Zenith	Seeded	Standard entry
6	PZA 32	Seeded	Patten Seed Company
7	PZB 33	Seeded	Patten Seed Company
8	Companion	Seeded	Seed Research of OR, Inc.
9	PST-R7ZM	Seeded	Pure-Seed Testing, Inc.
10	PST-R7MA	Seeded	Pure-Seed Testing, Inc.
11	GN-Z	Vegetative	Greg Norman Turf
12	DALZ 0102	Vegetative	Turfgrass America
13	DALZ 0104	Vegetative	Turfgrass America
14	BMZ 230	Vegetative	Turfgrass America
15	DALZ 9604	Vegetative	Turfgrass America
16	DALZ 0105	Vegetative	Turfgrass America
17	DALZ 0101	Vegetative	Turfgrass America
18	Zorro	Vegetative	Standard entry
19	6186	Vegetative	Bladerunner Farms
20	Chinese Common	Seeded	Standard entry

TABLE A.

2005 LOCATIONS, SITE DESCRIPTIONS AND MANAGEMENT PRACTICES IN  
THE 2002 NATIONAL ZOYSIAGRASS TEST

LOCATION	SOIL TEXTURE	SOIL PH	SOIL PHOSPHOROUS (LBS/ACRE)	SOIL POTASSIUM (LBS/ACRE)	NITROGEN (LBS/1000 SQ FT)	SUN OR SHADE	MOWING HEIGHT (IN)	IRRIGATION PRACTICED
AR1	SILT LOAM AND SILT	6.1-6.5	61-150	151-240	2.1-3.0	FULL SUN	0.6-1.0	TO PREVENT STRESS
CA3	SANDY LOAM	6.6-7.0	0-60	0-150	1.1-2.0	FULL SUN	1.6-2.0	TO PREVENT DORMANCY
FL3	-	-	-	-	-	-	-	-
GA1	SANDY LOAM	6.1-6.5	0-60	151-240	3.1-4.0	FULL SUN	0.6-1.0	TO PREVENT STRESS
IL2	SILTY CLAY LOAM	6.1-6.5	0-60	0-150	3.1-4.0	FULL SUN	0.6-1.0	TO PREVENT STRESS
IN1	SILT LOAM AND SILT	7.1-7.5	151-270	501+	1.1-2.0	FULL SUN	0.0-0.5	TO PREVENT DORMANCY
KS1	SILT LOAM AND SILT	6.6-7.0	151-270	241-375	3.1-4.0	FULL SUN	0.6-1.0	TO PREVENT STRESS
KY1	SILT LOAM AND SILT	6.1-6.5	61-150	241-375	2.1-3.0	FULL SUN	0.6-1.0	ONLY DURING SEVERE STRESS
MD1	SILT LOAM AND SILT	5.6-6.0	61-150	0-150	0.0-1.0	FULL SUN	0.0-0.5	TO PREVENT DORMANCY
MO1	SILT LOAM AND SILT	6.1-6.5	0-60	151-240	2.1-3.0	FULL SUN	0.6-1.0	TO PREVENT STRESS
MS1	SANDY LOAM	6.6-7.0	151-270	241-375	2.1-3.0	FULL SUN	1.6-2.0	TO PREVENT DORMANCY
NC1	SILTY CLAY AND CLAY	6.1-6.5	61-150	0-150	4.1-5.0	FULL SUN	0.6-1.0	TO PREVENT STRESS
NM1	SILTY CLAY LOAM	7.6-8.5	0-60	0-150	3.1-4.0	FULL SUN	1.6-2.0	TO PREVENT STRESS
OK1	SANDY LOAM	6.6-7.0	61-150	241-375	2.1-3.0	FULL SUN	0.0-0.5	TO PREVENT STRESS
SC1	SANDY LOAM	5.6-6.0	61-150	0-150	1.1-2.0	FULL SUN	2.6-3.0	TO PREVENT STRESS
TX1	SILTY CLAY AND CLAY	7.6-8.5	151-270	241-375	3.1-4.0	FULL SUN	0.6-1.0	TO PREVENT STRESS

TABLE B.

## LOCATIONS AND DATA COLLECTED IN 2005

LOCATION	JANUARY	FEBRUARY	MARCH	APRIL	MAY	JUNE	JULY	AUGUST	SEPTEMBER	OCTOBER	NOVEMBER	DECEMBER	GENETIC COLOR	SPRING GREENUP	LEAF TEXTURE
	QUALITY RATING														
AR1				X		X	X	X	X				X	X	X
CA3	X	X	X	X	X	X	X	X	X	X	X	X	X		
FL3				X	X		X	X	X	X	X	X	X		
GA1					X	X	X	X	X	X				X	
IL2						X	X	X	X	X	X			X	X
IN1						X	X	X	X	X	X			X	X
KS1							X	X							
KY1					X	X	X	X	X				X	X	
MD1					X	X	X	X	X					X	
MO1		X	X	X	X	X	X	X	X	X				X	X
MS1			X	X	X	X	X	X	X	X	X		X	X	X
NC1			X	X	X	X	X	X	X	X	X		X	X	X
NM1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
OK1					X	X	X	X	X	X	X		X	X	X
SC1				X	X	X	X	X	X	X	X		X	X	X
TX1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X

TABLE B. (CONT'D)

## LOCATIONS AND DATA COLLECTED IN 2005

LOCATION	SPRING DENSITY	SUMMER DENSITY	FALL DENSITY	PERCENT COVER SPRING	PERCENT COVER SUMMER	PERCENT COVER FALL	FROST TOLERANCE	WINTER COLOR	DOLLAR SPOT	BROWN PATCH WARM TEMP.	TAKE-ALL PATCH	MOLE CRICKET
AR1							X					
CA3								X				
FL3	X	X	X					X				
GA1												
IL2				X	X	X	X					
IN1												
KS1							X					
KY1												
MD1												
M01							X					
MS1		X										
NC1												
NM1	X	X	X					X	X			
OK1		X				X		X				
SC1							X		X	X		X
TX1	X	X	X								X	

TABLE B. (CONT'D)

## LOCATIONS AND DATA COLLECTED IN 2005

LOCATION	FALL COLOR SEPTEMBER	FALL COLOR OCTOBER	FALL COLOR NOVEMBER	FALL COLOR DECEMBER	SEEDHEAD RATINGS	MOWING QUALITY	MEAN ZOYSIA PATCH	PERCENT JUNE	ZOYSIA PATCH SEPTEMBER	LARGE PATCH RATINGS			
										MAY	JUNE	JULY	SEPTEMBER
AR1					X								
CA3					X								
FL3	X	X	X										
GA1	X	X											
IL2													
IN1								X	X				
KS1		X											
KY1													
MD1													
MO1		X											
MS1			X		X								
NC1			X								X	X	X
NM1	X	X	X	X	X	X							
OK1			X										
SC1		X			X								
TX1	X	X	X	X									

TABLE 1A.

MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS  
 GROWN AT ELEVEN LOCATIONS IN THE U.S. 1/  
 MAINTAINED USING "SCHEDULE A" \*\*  
 2005 DATA

NAME	TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/											
	AR1	GA1	IL2	IN1	KS1	KY1	MD1	MO1	NC1	OK1	TX1	MEAN
* ZORRO	7.0	7.9	8.1	6.3	5.2	7.6	7.4	5.0	8.0	7.8	6.6	7.0
* EMERALD	6.9	7.4	8.0	5.8	7.5	5.7	7.5	5.7	8.0	7.9	5.9	6.9
DALZ 0101	6.5	8.2	7.3	5.8	6.3	6.7	7.4	5.5	7.3	8.1	6.1	6.8
HIMENO	6.2	7.5	6.3	4.8	5.8	7.7	6.8	4.8	7.6	6.1	4.6	6.2
DALZ 0102	4.6	6.8	6.2	5.6	7.8	6.7	6.1	4.7	6.9	7.4	5.1	6.2
GN-Z	6.3	6.3	7.2	.	4.7	6.3	5.1	4.6	7.7	7.8	5.6	6.2
DALZ 0104	6.4	7.1	6.1	.	4.7	5.7	2.3	.	6.5	7.1	6.3	5.8
BMZ 230	5.2	5.8	5.5	4.9	6.3	7.4	4.2	5.3	7.8	6.4	4.4	5.8
* MEYER	5.9	6.6	5.9	6.3	4.3	8.1	4.5	3.3	6.3	7.2	3.0	5.6
DALZ 0105	5.3	7.5	4.7	.	4.5	5.4	2.4	.	5.9	6.6	6.0	5.4
PST-R7ZM	5.8	6.1	3.8	6.2	3.8	7.2	4.2	3.9	6.7	6.3	2.9	5.2
PST-R7MA	4.2	6.1	4.9	5.8	4.8	7.2	3.9	4.2	6.8	5.7	2.9	5.1
DALZ 9604	4.6	7.3	2.4	.	4.3	5.4	2.1	.	7.2	7.2	5.4	5.1
* ZENITH	4.5	6.1	5.2	5.0	4.8	6.3	3.5	4.5	7.3	5.2	3.4	5.1
PZB 33	4.4	6.2	5.1	4.6	5.0	6.2	3.7	5.0	7.2	5.0	3.4	5.1
PZA 32	3.1	6.1	5.2	5.5	5.2	6.5	3.8	4.4	6.8	4.9	3.1	5.0
6186	3.7	6.4	1.4	.	4.0	6.7	2.8	3.6	6.4	7.2	5.4	4.8
* COMPANION	4.5	5.9	4.3	4.0	4.8	5.7	3.3	3.7	6.5	4.9	2.9	4.6
* CHINESE COMMON	3.7	5.8	4.0	3.9	5.2	6.4	2.9	4.8	6.7	3.8	3.2	4.6
* J-37	4.6	5.2	4.3	4.3	4.8	6.2	3.0	4.8	6.1	3.8	2.7	4.5
LSD VALUE	0.9	0.6	1.3	1.3	1.5	1.1	1.0	1.1	0.5	1.3	0.6	0.3
C.V. (%)	11.0	5.6	15.4	15.0	17.6	10.7	13.1	13.5	4.8	12.7	8.2	11.7

\* COMMERCIALLY AVAILABLE IN THE USA IN 2006.

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

\*\* SCHEDULE A - 0.5 -0.75 INCH MOWING HEIGHT  
 0.5 -0.66 lbs. N/1000 FT2/GROWING MONTH  
 IRRIGATION TO PREVENT STRESS  
 PEST CONTROL - PRE-EMERGENT ANNUAL GRASS CONTROL ALLOWED, BROADLEAF WEED CONTROL AS NEEDED.  
 INSECTICIDE AND FUNGICIDE APPLICATIONS MAY BE APPLIED ON A CURATIVE BASIS ONLY AFTER DATA ON CULTIVAR RESPONSE IS COLLECTED.

TABLE 1B.

MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS  
 GROWN AT ELEVEN LOCATIONS IN THE U.S. 1/  
 MAINTAINED USING "SCHEDULE A" \*

2005 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

NAME	AR1	GA1	IL2	IN1	KS1	KY1	MD1	MO1	NC1	OK1	TX1	MEAN
PST-R7ZM	5.8	6.1	3.8	6.2	3.8	7.2	4.2	3.9	6.7	6.3	2.9	5.2
PST-R7MA	4.2	6.1	4.9	5.8	4.8	7.2	3.9	4.2	6.8	5.7	2.9	5.1
ZENITH	4.5	6.1	5.2	5.0	4.8	6.3	3.5	4.5	7.3	5.2	3.4	5.1
PZB 33	4.4	6.2	5.1	4.6	5.0	6.2	3.7	5.0	7.2	5.0	3.4	5.1
PZA 32	3.1	6.1	5.2	5.5	5.2	6.5	3.8	4.4	6.8	4.9	3.1	5.0
COMPANION	4.5	5.9	4.3	4.0	4.8	5.7	3.3	3.7	6.5	4.9	2.9	4.6
CHINESE COMMON	3.7	5.8	4.0	3.9	5.2	6.4	2.9	4.8	6.7	3.8	3.2	4.6
J-37	4.6	5.2	4.3	4.3	4.8	6.2	3.0	4.8	6.1	3.8	2.7	4.5
LSD VALUE	1.0	0.6	0.9	1.2	1.5	1.1	1.2	1.2	0.4	1.6	0.6	0.3
C.V. (%)	14.1	6.5	12.0	15.7	19.9	10.7	20.2	16.5	3.9	20.4	13.0	13.9

TABLE 1C.

MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS  
 GROWN AT ELEVEN LOCATIONS IN THE U.S. 1/  
 MAINTAINED USING "SCHEDULE A" \*

2005 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

NAME	AR1	GA1	IL2	IN1	KS1	KY1	MD1	MO1	NC1	OK1	TX1	MEAN
ZORRO	7.0	7.9	8.1	6.3	5.2	7.6	7.4	5.0	8.0	7.8	6.6	7.0
EMERALD	6.9	7.4	8.0	5.8	7.5	5.7	7.5	5.7	8.0	7.9	5.9	6.9
DALZ 0101	6.5	8.2	7.3	5.8	6.3	6.7	7.4	5.5	7.3	8.1	6.1	6.8
HIMENO	6.2	7.5	6.3	4.8	5.8	7.7	6.8	4.8	7.6	6.1	4.6	6.2
DALZ 0102	4.6	6.8	6.2	5.6	7.8	6.7	6.1	4.7	6.9	7.4	5.1	6.2
GN-Z	6.3	6.3	7.2	.	4.7	6.3	5.1	4.6	7.7	7.8	5.6	6.2
DALZ 0104	6.4	7.1	6.1	.	4.7	5.7	2.3	.	6.5	7.1	6.3	5.8
BMZ 230	5.2	5.8	5.5	4.9	6.3	7.4	4.2	5.3	7.8	6.4	4.4	5.8
MEYER	5.9	6.6	5.9	6.3	4.3	8.1	4.5	3.3	6.3	7.2	3.0	5.6
DALZ 0105	5.3	7.5	4.7	.	4.5	5.4	2.4	.	5.9	6.6	6.0	5.4
DALZ 9604	4.6	7.3	2.4	.	4.3	5.4	2.1	.	7.2	7.2	5.4	5.1
6186	3.7	6.4	1.4	.	4.0	6.7	2.8	3.6	6.4	7.2	5.4	4.8
LSD VALUE	0.9	0.6	1.5	1.3	1.4	1.2	0.8	0.9	0.6	1.0	0.6	0.3
C.V. (%)	9.4	5.1	16.6	14.3	16.3	10.6	8.9	9.6	5.3	8.7	6.4	10.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
 STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

\*\* SCHEDULE A - 0.5 - 0.75 INCH MOWING HEIGHT  
 0.5 - 0.66 lbs. N/1000 FT<sup>2</sup>/GROWING MONTH  
 IRRIGATION TO PREVENT STRESS

PEST CONTROL - PRE-EMERGENT ANNUAL GRASS CONTROL ALLOWED, BROADLEAF WEED CONTROL AS NEEDED.  
 INSECTICIDE AND FUNGICIDE APPLICATIONS MAY BE APPLIED ON A CURATIVE BASIS ONLY  
 AFTER DATA ON CULTIVAR RESPONSE IS COLLECTED.

TABLE 2A. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS  
 GROWN AT FIVE LOCATIONS IN THE U.S. 1/  
 MAINTAINED USING "SCHEDULE B" \*  
 2005 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

NAME	CA3	FL3	MS1	NM1	SC1	MEAN
ZORRO	5.2	7.8	7.2	5.1	8.2	6.7
EMERALD	5.2	7.5	7.4	5.3	7.8	6.6
DALZ 0101	5.3	7.7	7.0	5.4	6.8	6.4
GN-Z	5.4	7.6	6.5	5.1	7.3	6.4
DALZ 0104	4.8	7.3	6.3	4.8	7.3	6.1
DALZ 0102	5.0	7.6	6.2	4.5	6.9	6.0
BMZ 230	5.0	7.3	5.7	4.5	7.5	6.0
6186	4.9	7.0	5.5	5.1	7.0	5.9
DALZ 9604	4.9	7.1	5.3	4.9	7.1	5.9
DALZ 0105	4.9	6.3	6.2	4.9	6.6	5.8
ZENITH	5.1	6.8	5.7	4.2	7.0	5.8
PZB 33	4.8	6.7	5.4	4.2	7.3	5.7
HIMENO	3.0	7.0	5.7	4.1	8.0	5.6
MEYER	4.2	6.2	6.5	3.7	6.9	5.5
PZA 32	4.6	6.5	5.0	4.2	6.8	5.4
PST-R7MA	3.8	6.2	5.9	4.1	6.9	5.4
J-37	4.5	7.0	4.6	4.2	6.3	5.3
PST-R7ZM	4.0	6.1	5.8	3.5	7.0	5.3
COMPANION	4.6	6.1	5.4	4.1	5.9	5.2
CHINESE COMMON	4.6	6.4	4.9	4.1	6.0	5.2
LSD VALUE	0.5	0.7	0.4	0.4	1.3	0.3
C.V. (%)	5.9	6.0	4.7	5.7	11.5	7.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
 STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

\* SCHEDULE B - 1.5 - 2 INCH MOWING HEIGHT WITH ROTARY MOWER  
 0.25 - 0.33 lbs. N/1000 FT<sup>2</sup>/GROWING MONTH  
 IRRIGATION TO PREVENT DORMANCY OR SEVERE STRESS  
 PEST CONTROL - PRE-EMERGENT ANNUAL GRASS CONTROL ALLOWED, BROADLEAF WEED CONTROL AS NEEDED.  
 INSECTICIDE AND FUNGICIDE APPLICATIONS MAY BE APPLIED ON A CURATIVE BASIS ONLY  
 AFTER DATA ON CULTIVAR RESPONSE IS COLLECTED.

TABLE 2B. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS  
GROWN AT FIVE LOCATIONS IN THE U.S. 1/  
MAINTAINED USING "SCHEDULE B" \*  
2005 DATA

NAME	CA3	FL3	MS1	NM1	SC1	MEAN
ZENITH	5.1	6.8	5.7	4.2	7.0	5.8
PZB 33	4.8	6.7	5.4	4.2	7.3	5.7
PZA 32	4.6	6.5	5.0	4.2	6.8	5.4
PST-R7MA	3.8	6.2	5.9	4.1	6.9	5.4
J-37	4.5	7.0	4.6	4.2	6.3	5.3
PST-R7ZM	4.0	6.1	5.8	3.5	7.0	5.3
COMPANION	4.6	6.1	5.4	4.1	5.9	5.2
CHINESE COMMON	4.6	6.4	4.9	4.1	6.0	5.2
LSD VALUE	0.6	0.7	0.4	0.3	1.6	0.4
C.V. (%)	7.6	7.0	4.9	5.3	14.7	9.9

TABLE 2C. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS  
GROWN AT FIVE LOCATIONS IN THE U.S. 1/  
MAINTAINED USING "SCHEDULE B" \*  
2005 DATA

NAME	CA3	FL3	MS1	NM1	SC1	MEAN
ZORRO	5.2	7.8	7.2	5.1	8.2	6.7
EMERALD	5.2	7.5	7.4	5.3	7.8	6.6
DALZ 0101	5.3	7.7	7.0	5.4	6.8	6.4
GN-Z	5.4	7.6	6.5	5.1	7.3	6.4
DALZ 0104	4.8	7.3	6.3	4.8	7.3	6.1
DALZ 0102	5.0	7.6	6.2	4.5	6.9	6.0
BMZ 230	5.0	7.3	5.7	4.5	7.5	6.0
6186	4.9	7.0	5.5	5.1	7.0	5.9
DALZ 9604	4.9	7.1	5.3	4.9	7.1	5.9
DALZ 0105	4.9	6.3	6.2	4.9	6.6	5.8
HIMENO	3.0	7.0	5.7	4.1	8.0	5.6
MEYER	4.2	6.2	6.5	3.7	6.9	5.5
LSD VALUE	0.4	0.6	0.5	0.4	1.1	0.3
C.V. (%)	4.6	5.3	4.5	5.8	9.1	6.6

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

\* SCHEDULE B - 1.5 - 2 INCH MOWING HEIGHT WITH ROTARY MOWER  
0.25 - 0.33 lbs. N/1000 FT<sup>2</sup>/GROWING MONTH  
IRRIGATION TO PREVENT DORMANCY OR SEVERE STRESS  
PEST CONTROL - PRE-EMERGENT ANNUAL GRASS CONTROL ALLOWED, BROADLEAF WEED CONTROL AS NEEDED.  
INSECTICIDE AND FUNGICIDE APPLICATIONS MAY BE APPLIED ON A CURATIVE BASIS ONLY  
AFTER DATA ON CULTIVAR RESPONSE IS COLLECTED.

TABLE 3A. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS  
 GROWN AT EIGHT LOCATIONS 1/  
 IN THE TRANSITION REGION  
 2005 DATA

NAME	TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/								MEAN
	AR1	IL2	KS1	KY1	MD1	MO1	NC1	OK1	
EMERALD	6.9	8.0	7.5	5.7	7.5	5.7	8.0	7.9	7.1
ZORRO	7.0	8.1	5.2	7.6	7.4	5.0	8.0	7.8	7.0
DALZ 0101	6.5	7.3	6.3	6.7	7.4	5.5	7.3	8.1	6.9
HIMENO	6.2	6.3	5.8	7.7	6.8	4.8	7.6	6.1	6.4
DALZ 0102	4.6	6.2	7.8	6.7	6.1	4.7	6.9	7.4	6.3
GN-Z	6.3	7.2	4.7	6.3	5.1	4.6	7.7	7.8	6.2
BMZ 230	5.2	5.5	6.3	7.4	4.2	5.3	7.8	6.4	6.0
MEYER	5.9	5.9	4.3	8.1	4.5	3.3	6.3	7.2	5.7
DALZ 0104	6.4	6.1	4.7	5.7	2.3	.	6.5	7.1	5.5
PST-R7MA	4.2	4.9	4.8	7.2	3.9	4.2	6.8	5.7	5.2
PST-R7ZM	5.8	3.8	3.8	7.2	4.2	3.9	6.7	6.3	5.2
PZB 33	4.4	5.1	5.0	6.2	3.7	5.0	7.2	5.0	5.2
ZENITH	4.5	5.2	4.8	6.3	3.5	4.5	7.3	5.2	5.2
PZA 32	3.1	5.2	5.2	6.5	3.8	4.4	6.8	4.9	5.0
DALZ 0105	5.3	4.7	4.5	5.4	2.4	.	5.9	6.6	5.0
DALZ 9604	4.6	2.4	4.3	5.4	2.1	.	7.2	7.2	4.8
COMPANION	4.5	4.3	4.8	5.7	3.3	3.7	6.5	4.9	4.7
J-37	4.6	4.3	4.8	6.2	3.0	4.8	6.1	3.8	4.7
CHINESE COMMON	3.7	4.0	5.2	6.4	2.9	4.8	6.7	3.8	4.7
6186	3.7	1.4	4.0	6.7	2.8	3.6	6.4	7.2	4.5
LSD VALUE	0.9	1.3	1.5	1.1	1.0	1.1	0.5	1.3	0.4
C.V. (%)	11.0	15.4	17.6	10.7	13.1	13.5	4.8	12.7	12.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
 STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 3B. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS  
 GROWN AT EIGHT LOCATIONS 1/  
 IN THE TRANSITION REGION  
 2005 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/									
NAME	AR1	IL2	KS1	KY1	MD1	MO1	NC1	OK1	MEAN
PST-R7MA	4.2	4.9	4.8	7.2	3.9	4.2	6.8	5.7	5.2
PST-R7ZM	5.8	3.8	3.8	7.2	4.2	3.9	6.7	6.3	5.2
PZB 33	4.4	5.1	5.0	6.2	3.7	5.0	7.2	5.0	5.2
ZENITH	4.5	5.2	4.8	6.3	3.5	4.5	7.3	5.2	5.2
PZA 32	3.1	5.2	5.2	6.5	3.8	4.4	6.8	4.9	5.0
COMPANION	4.5	4.3	4.8	5.7	3.3	3.7	6.5	4.9	4.7
J-37	4.6	4.3	4.8	6.2	3.0	4.8	6.1	3.8	4.7
CHINESE COMMON	3.7	4.0	5.2	6.4	2.9	4.8	6.7	3.8	4.7
LSD VALUE	1.0	0.9	1.5	1.1	1.2	1.2	0.4	1.6	0.4
C.V. (%)	14.1	12.0	19.9	10.7	20.2	16.5	3.9	20.4	14.6

TABLE 3C. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS  
 GROWN AT EIGHT LOCATIONS 1/  
 IN THE TRANSITION REGION  
 2005 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/									
NAME	AR1	IL2	KS1	KY1	MD1	MO1	NC1	OK1	MEAN
EMERALD	6.9	8.0	7.5	5.7	7.5	5.7	8.0	7.9	7.1
ZORRO	7.0	8.1	5.2	7.6	7.4	5.0	8.0	7.8	7.0
DALZ 0101	6.5	7.3	6.3	6.7	7.4	5.5	7.3	8.1	6.9
HIMENO	6.2	6.3	5.8	7.7	6.8	4.8	7.6	6.1	6.4
DALZ 0102	4.6	6.2	7.8	6.7	6.1	4.7	6.9	7.4	6.3
GN-Z	6.3	7.2	4.7	6.3	5.1	4.6	7.7	7.8	6.2
BMZ 230	5.2	5.5	6.3	7.4	4.2	5.3	7.8	6.4	6.0
MEYER	5.9	5.9	4.3	8.1	4.5	3.3	6.3	7.2	5.7
DALZ 0104	6.4	6.1	4.7	5.7	2.3	.	6.5	7.1	5.5
DALZ 0105	5.3	4.7	4.5	5.4	2.4	.	5.9	6.6	5.0
DALZ 9604	4.6	2.4	4.3	5.4	2.1	.	7.2	7.2	4.8
6186	3.7	1.4	4.0	6.7	2.8	3.6	6.4	7.2	4.5
LSD VALUE	0.9	1.5	1.4	1.2	0.8	0.9	0.6	1.0	0.4
C.V. (%)	9.4	16.6	16.3	10.6	8.9	9.6	5.3	8.7	11.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
 STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 4A. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS  
 GROWN AT FIVE LOCATIONS 1/  
 IN THE SOUTHEAST REGION  
 2005 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

NAME	FL3	GA1	MS1	SC1	TX1	MEAN
ZORRO	7.8	7.9	7.2	8.2	6.6	7.5
EMERALD	7.5	7.4	7.4	7.8	5.9	7.2
DALZ 0101	7.7	8.2	7.0	6.8	6.1	7.1
DALZ 0104	7.3	7.1	6.3	7.3	6.3	6.9
GN-Z	7.6	6.3	6.5	7.3	5.6	6.7
HIMENO	7.0	7.5	5.7	8.0	4.6	6.6
DALZ 0105	6.3	7.5	6.2	6.6	6.0	6.5
DALZ 0102	7.6	6.8	6.2	6.9	5.1	6.5
DALZ 9604	7.1	7.3	5.3	7.1	5.4	6.5
6186	7.0	6.4	5.5	7.0	5.4	6.3
BMZ 230	7.3	5.8	5.7	7.5	4.4	6.2
MEYER	6.2	6.6	6.5	6.9	3.0	5.8
ZENITH	6.8	6.1	5.7	7.0	3.4	5.8
PZB 33	6.7	6.2	5.4	7.3	3.4	5.8
PST-R7ZM	6.1	6.1	5.8	7.0	2.9	5.6
PST-R7MA	6.2	6.1	5.9	6.9	2.9	5.6
PZA 32	6.5	6.1	5.0	6.8	3.1	5.5
CHINESE COMMON	6.4	5.8	4.9	6.0	3.2	5.3
COMPANION	6.1	5.9	5.4	5.9	2.9	5.2
J-37	7.0	5.2	4.6	6.3	2.7	5.2
LSD VALUE	0.7	0.6	0.4	1.3	0.6	0.3
C.V. (%)	6.0	5.6	4.7	11.5	8.2	7.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
 STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 4B. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS  
 GROWN AT FIVE LOCATIONS 1/  
 IN THE SOUTHEAST REGION  
 2005 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

NAME	FL3	GA1	MS1	SC1	TX1	MEAN
ZENITH	6.8	6.1	5.7	7.0	3.4	5.8
PZB 33	6.7	6.2	5.4	7.3	3.4	5.8
PST-R7ZM	6.1	6.1	5.8	7.0	2.9	5.6
PST-R7MA	6.2	6.1	5.9	6.9	2.9	5.6
PZA 32	6.5	6.1	5.0	6.8	3.1	5.5
CHINESE COMMON	6.4	5.8	4.9	6.0	3.2	5.3
COMPANION	6.1	5.9	5.4	5.9	2.9	5.2
J-37	7.0	5.2	4.6	6.3	2.7	5.2
LSD VALUE	0.7	0.6	0.4	1.6	0.6	0.4
C.V. (%)	7.0	6.5	4.9	14.7	13.0	10.1

TABLE 4C. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS  
 GROWN AT FIVE LOCATIONS 1/  
 IN THE SOUTHEAST REGION  
 2005 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

NAME	FL3	GA1	MS1	SC1	TX1	MEAN
ZORRO	7.8	7.9	7.2	8.2	6.6	7.5
EMERALD	7.5	7.4	7.4	7.8	5.9	7.2
DALZ 0101	7.7	8.2	7.0	6.8	6.1	7.1
DALZ 0104	7.3	7.1	6.3	7.3	6.3	6.9
GN-Z	7.6	6.3	6.5	7.3	5.6	6.7
HIMENO	7.0	7.5	5.7	8.0	4.6	6.6
DALZ 0105	6.3	7.5	6.2	6.6	6.0	6.5
DALZ 0102	7.6	6.8	6.2	6.9	5.1	6.5
DALZ 9604	7.1	7.3	5.3	7.1	5.4	6.5
6186	7.0	6.4	5.5	7.0	5.4	6.3
BMZ 230	7.3	5.8	5.7	7.5	4.4	6.2
MEYER	6.2	6.6	6.5	6.9	3.0	5.8
LSD VALUE	0.6	0.6	0.5	1.1	0.6	0.3
C.V. (%)	5.3	5.1	4.5	9.1	6.4	6.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 5A. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS FOR EACH MONTH GROWN AT WEST LAFAYETTE, IN (NORTH CENTRAL) 1/  
2005 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF: MONTHS 2/

NAME	MAY	JUN	JUL	AUG	SEP	OCT	MEAN
MEYER	6.3	7.7	7.7	6.3	4.0	5.7	6.3
ZORRO	3.7	7.3	6.3	7.7	5.0	7.7	6.3
PST-R7ZM	7.3	7.3	5.7	6.0	5.7	5.3	6.2
DALZ 0101	3.3	6.7	7.3	8.0	4.0	5.7	5.8
EMERALD	3.7	6.7	7.3	8.0	3.3	6.0	5.8
PST-R7MA	6.3	7.0	4.7	5.7	6.3	4.7	5.8
DALZ 0102	6.7	7.3	5.0	4.7	4.0	5.7	5.6
PZA 32	7.3	6.0	5.0	4.0	5.3	5.3	5.5
ZENITH	6.7	6.3	4.3	3.7	4.3	4.7	5.0
BMZ 230	7.3	6.3	3.3	3.3	5.0	4.0	4.9
HIMENO	7.0	6.3	4.0	3.3	4.3	3.7	4.8
PZB 33	6.0	4.7	3.7	4.0	4.3	4.7	4.6
J-37	5.3	5.0	3.0	3.0	5.0	4.7	4.3
COMPANION	5.3	5.0	3.3	2.7	4.3	3.3	4.0
CHINESE COMMON	5.0	4.0	3.3	2.7	5.3	3.3	3.9
6186	.	.	.	.	.	.	.
DALZ 0104	.	.	.	.	.	.	.
DALZ 0105	.	.	.	.	.	.	.
DALZ 9604	.	.	.	.	.	.	.
GN-Z	.	.	.	.	.	.	.
LSD VALUE	2.2	3.0	1.9	1.8	2.1	2.5	1.4
C.V. (%)	23.8	29.5	23.4	22.5	28.3	30.9	16.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 5B. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS FOR EACH MONTH GROWN AT WEST LAFAYETTE, IN (NORTH CENTRAL) 1/  
2005 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF: MONTHS 2/

NAME	MAY	JUN	JUL	AUG	SEP	OCT	MEAN
PST-R7ZM	7.3	7.3	5.7	6.0	5.7	5.3	6.2
PST-R7MA	6.3	7.0	4.7	5.7	6.3	4.7	5.8
PZA 32	7.3	6.0	5.0	4.0	5.3	5.3	5.5
ZENITH	6.7	6.3	4.3	3.7	4.3	4.7	5.0
PZB 33	6.0	4.7	3.7	4.0	4.3	4.7	4.6
J-37	5.3	5.0	3.0	3.0	5.0	4.7	4.3
COMPANION	5.3	5.0	3.3	2.7	4.3	3.3	4.0
CHINESE COMMON	5.0	4.0	3.3	2.7	5.3	3.3	3.9
LSD VALUE	1.9	2.7	1.6	1.2	1.7	1.9	1.2
C.V. (%)	19.0	29.3	24.2	18.6	21.2	26.8	15.7

TABLE 5C. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS FOR EACH MONTH GROWN AT WEST LAFAYETTE, IN (NORTH CENTRAL) 1/  
2005 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF: MONTHS 2/

NAME	MAY	JUN	JUL	AUG	SEP	OCT	MEAN
MEYER	6.3	7.7	7.7	6.3	4.0	5.7	6.3
ZORRO	3.7	7.3	6.3	7.7	5.0	7.7	6.3
DALZ 0101	3.3	6.7	7.3	8.0	4.0	5.7	5.8
EMERALD	3.7	6.7	7.3	8.0	3.3	6.0	5.8
DALZ 0102	6.7	7.3	5.0	4.7	4.0	5.7	5.6
BMZ 230	7.3	6.3	3.3	3.3	5.0	4.0	4.9
HIMENO	7.0	6.3	4.0	3.3	4.3	3.7	4.8
6186	.	.	.	.	.	.	.
DALZ 0104	.	.	.	.	.	.	.
DALZ 0105	.	.	.	.	.	.	.
DALZ 9604	.	.	.	.	.	.	.
GN-Z	.	.	.	.	.	.	.
LSD VALUE	2.7	3.4	2.2	2.5	2.7	3.2	1.6
C.V. (%)	31.3	30.8	23.7	26.1	39.6	36.2	17.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 6A. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS  
 GROWN AT TWO LOCATIONS 1/  
 IN THE SOUTHWEST REGION  
 2005 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

NAME	CA3	NM1	MEAN
DALZ 0101	5.3	5.4	5.4
EMERALD	5.2	5.3	5.3
GN-Z	5.4	5.1	5.3
ZORRO	5.2	5.1	5.2
6186	4.9	5.1	5.0
DALZ 9604	4.9	4.9	4.9
DALZ 0105	4.9	4.9	4.9
DALZ 0104	4.8	4.8	4.8
BMZ 230	5.0	4.5	4.8
DALZ 0102	5.0	4.5	4.8
ZENITH	5.1	4.2	4.6
PZB 33	4.8	4.2	4.5
PZA 32	4.6	4.2	4.4
J-37	4.5	4.2	4.3
COMPANION	4.6	4.1	4.3
CHINESE COMMON	4.6	4.1	4.3
MEYER	4.2	3.7	4.0
PST-R7MA	3.8	4.1	3.9
PST-R7ZM	4.0	3.5	3.7
HIMENO	3.0	4.1	3.6
LSD VALUE	0.5	0.4	0.3
C.V. (%)	5.9	5.7	5.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
 STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 6B. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS  
 GROWN AT TWO LOCATIONS 1/  
 IN THE SOUTHWEST REGION  
 2005 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

NAME	CA3	NM1	MEAN
ZENITH	5.1	4.2	4.6
PZB 33	4.8	4.2	4.5
PZA 32	4.6	4.2	4.4
J-37	4.5	4.2	4.3
COMPANION	4.6	4.1	4.3
CHINESE COMMON	4.6	4.1	4.3
PST-R7MA	3.8	4.1	3.9
PST-R7ZM	4.0	3.5	3.7
LSD VALUE	0.6	0.3	0.3
C.V. (%)	7.6	5.3	6.5

TABLE 6C. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS  
 GROWN AT TWO LOCATIONS 1/  
 IN THE SOUTHWEST REGION  
 2005 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

NAME	CA3	NM1	MEAN
DALZ 0101	5.3	5.4	5.4
EMERALD	5.2	5.3	5.3
GN-Z	5.4	5.1	5.3
ZORRO	5.2	5.1	5.2
6186	4.9	5.1	5.0
DALZ 9604	4.9	4.9	4.9
DALZ 0105	4.9	4.9	4.9
DALZ 0104	4.8	4.8	4.8
BMZ 230	5.0	4.5	4.8
DALZ 0102	5.0	4.5	4.8
MEYER	4.2	3.7	4.0
HIMENO	3.0	4.1	3.6
LSD VALUE	0.4	0.4	0.3
C.V. (%)	4.6	5.8	5.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
 STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 7A.

GENETIC COLOR RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2005 DATA

GENETIC COLOR RATINGS 1-9; 9=DARK GREEN 2/

NAME	AR1	CA3	FL3	GA1	IL2	IN1	KY1	MS1	NC1	NM1	OK1	SC1	TX1	MEAN
HIMENO	6.0	.	6.7	8.7	6.7	6.3	8.3	7.0	8.0	6.0	9.0	7.3	9.0	7.4
EMERALD	6.7	7.0	7.7	8.3	7.3	5.3	6.7	6.7	7.7	6.7	7.0	7.3	7.3	7.1
MEYER	6.7	6.7	5.0	9.0	7.7	5.7	7.7	6.0	8.0	5.7	9.0	6.3	7.7	7.0
ZENITH	6.0	7.0	6.7	7.7	5.7	7.3	7.0	6.0	8.0	5.0	8.0	6.7	8.0	6.8
6186	7.0	5.3	6.3	9.0	9.0	5.0	8.5	6.0	8.0	5.0	8.0	4.0	7.0	6.8
DALZ 0104	6.7	6.3	6.3	8.0	6.7	.	8.3	6.0	7.7	6.0	8.0	4.7	6.7	6.8
DALZ 0101	5.3	7.0	7.7	8.3	6.3	7.7	5.0	6.0	6.7	6.3	6.7	6.7	7.3	6.7
ZORRO	7.0	6.0	7.3	7.3	5.0	5.7	7.3	6.0	7.7	6.3	6.3	7.7	7.0	6.7
PZA 32	4.0	7.0	5.7	7.7	6.3	5.3	7.0	6.7	8.0	5.7	7.3	6.7	7.7	6.5
PZB 33	6.0	7.0	5.7	7.0	6.3	5.3	6.7	6.0	8.0	5.3	7.7	6.0	7.3	6.5
GN-Z	5.3	6.7	6.7	7.7	5.0	.	7.7	6.0	6.7	6.0	6.0	6.7	7.3	6.5
COMPANION	5.3	7.0	5.7	7.3	5.7	6.3	6.7	6.0	7.7	5.3	7.7	5.3	7.7	6.4
PST-R7ZM	6.0	7.0	5.0	7.7	6.0	5.3	7.0	6.0	8.0	4.0	7.3	7.0	6.7	6.4
DALZ 9604	6.0	5.0	6.3	7.7	7.7	.	7.5	6.0	7.3	5.3	6.7	5.0	5.3	6.3
DALZ 0105	7.0	5.0	7.0	7.0	6.0	.	.	6.0	7.3	5.0	7.3	5.3	6.0	6.3
PST-R7MA	5.0	6.0	5.0	7.3	7.0	5.7	6.7	6.0	8.3	4.7	7.0	5.3	7.0	6.2
J-37	4.7	7.0	6.3	7.0	4.0	5.0	6.3	6.0	7.7	4.3	8.0	6.0	7.0	6.1
BMZ 230	5.0	7.0	6.7	7.7	2.3	5.0	6.3	6.0	7.3	5.0	7.3	7.0	6.0	6.1
CHINESE COMMON	3.0	7.0	6.0	7.5	2.3	5.7	6.7	6.0	7.3	5.0	7.3	4.7	7.0	5.8
DALZ 0102	6.0	6.7	7.0	7.3	1.3	2.7	5.3	6.0	6.7	4.3	6.0	6.3	6.3	5.5
LSD VALUE	1.7	0.7	0.9	0.9	1.8	1.6	2.1	0.3	0.9	1.5	0.7	1.9	1.4	0.4
C.V. (%)	18.0	6.0	8.4	7.0	19.7	16.9	17.8	3.0	7.4	17.1	6.3	19.6	12.5	12.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 7B.

GENETIC COLOR RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

## GENETIC COLOR RATINGS 1-9; 9=DARK GREEN 2/

NAME	AR1	CA3	FL3	GA1	IL2	IN1	KY1	MS1	NC1	NM1	OK1	SC1	TX1	MEAN
ZENITH	6.0	7	6.7	7.7	5.7	7.3	7.0	6.0	8.0	5.0	8.0	6.7	8.0	6.8
PZA 32	4.0	7	5.7	7.7	6.3	5.3	7.0	6.7	8.0	5.7	7.3	6.7	7.7	6.5
PZB 33	6.0	7	5.7	7.0	6.3	5.3	6.7	6.0	8.0	5.3	7.7	6.0	7.3	6.5
COMPANION	5.3	7	5.7	7.3	5.7	6.3	6.7	6.0	7.7	5.3	7.7	5.3	7.7	6.4
PST-R7ZM	6.0	7	5.0	7.7	6.0	5.3	7.0	6.0	8.0	4.0	7.3	7.0	6.7	6.4
PST-R7MA	5.0	6	5.0	7.3	7.0	5.7	6.7	6.0	8.3	4.7	7.0	5.3	7.0	6.2
J-37	4.7	7	6.3	7.0	4.0	5.0	6.3	6.0	7.7	4.3	8.0	6.0	7.0	6.1
CHINESE COMMON	3.0	7	6.0	7.5	2.3	5.7	6.7	6.0	7.3	5.0	7.3	4.7	7.0	5.8
LSD VALUE	2.0	0	0.9	0.9	1.9	1.3	1.2	0.3	0.7	1.4	0.9	2.5	1.3	0.4
C.V. (%)	25.2	0	10.0	7.1	21.3	14.2	11.3	3.4	5.2	17.1	7.7	26.3	11.2	13.3

TABLE 7C.

GENETIC COLOR RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

## GENETIC COLOR RATINGS 1-9; 9=DARK GREEN 2/

NAME	AR1	CA3	FL3	GA1	IL2	IN1	KY1	MS1	NC1	NM1	OK1	SC1	TX1	MEAN
HIMENO	6.0	.	6.7	8.7	6.7	6.3	8.3	7.0	8.0	6.0	9.0	7.3	9.0	7.4
EMERALD	6.7	7.0	7.7	8.3	7.3	5.3	6.7	6.7	7.7	6.7	7.0	7.3	7.3	7.1
MEYER	6.7	6.7	5.0	9.0	7.7	5.7	7.7	6.0	8.0	5.7	9.0	6.3	7.7	7.0
6186	7.0	5.3	6.3	9.0	9.0	5.0	8.5	6.0	8.0	5.0	8.0	4.0	7.0	6.8
DALZ 0104	6.7	6.3	6.3	8.0	6.7	.	8.3	6.0	7.7	6.0	8.0	4.7	6.7	6.8
DALZ 0101	5.3	7.0	7.7	8.3	6.3	7.7	5.0	6.0	6.7	6.3	6.7	6.7	7.3	6.7
ZORRO	7.0	6.0	7.3	7.3	5.0	5.7	7.3	6.0	7.7	6.3	6.3	7.7	7.0	6.7
GN-Z	5.3	6.7	6.7	7.7	5.0	.	7.7	6.0	6.7	6.0	6.0	6.7	7.3	6.5
DALZ 9604	6.0	5.0	6.3	7.7	7.7	.	7.5	6.0	7.3	5.3	6.7	5.0	5.3	6.3
DALZ 0105	7.0	5.0	7.0	7.0	6.0	.	.	6.0	7.3	5.0	7.3	5.3	6.0	6.3
BMZ 230	5.0	7.0	6.7	7.7	2.3	5.0	6.3	6.0	7.3	5.0	7.3	7.0	6.0	6.1
DALZ 0102	6.0	6.7	7.0	7.3	1.3	2.7	5.3	6.0	6.7	4.3	6.0	6.3	6.3	5.5
LSD VALUE	1.4	0.8	0.8	0.9	1.8	1.9	2.6	0.3	1.0	1.5	0.6	1.4	1.5	0.4
C.V. (%)	13.7	7.9	7.4	6.9	18.7	19.7	21.3	2.7	8.7	17.0	5.1	14.0	13.4	12.6

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 8A.

SPRING GREENUP RATINGS OF ZOysiagrass CULTIVARS 1/  
2005 DATA

## SPRING GREENUP RATINGS 1-9; 9=COMPLETELY GREEN 2/

NAME	AR1	IL2	IN1	KY1	MD1	MO1	MS1	NC1	NM1	OK1	SC1	TX1	MEAN
BMZ 230	8.0	5.3	6.3	9.0	6.0	4.0	7.0	8.0	2.7	6.0	7.3	4.7	6.2
ZENITH	6.3	8.0	5.3	5.7	5.0	4.3	5.7	7.3	3.0	3.7	6.3	3.7	5.4
PZB 33	7.3	8.3	3.7	6.0	4.0	4.0	5.7	7.7	3.0	4.0	5.7	4.3	5.3
MEYER	7.7	7.7	3.3	8.7	4.3	2.3	6.3	7.3	3.0	2.0	6.7	2.3	5.1
ZORRO	5.0	3.3	3.7	5.3	6.3	1.0	6.3	8.7	3.0	4.0	7.3	6.3	5.0
PST-R7ZM	7.3	6.3	4.3	6.0	4.0	2.3	5.7	7.7	3.3	2.3	6.7	4.0	5.0
DALZ 0102	2.3	5.0	4.0	8.7	6.0	1.3	5.7	7.7	3.0	6.0	5.7	3.7	4.9
PST-R7MA	6.3	6.3	5.0	6.3	3.3	3.7	5.0	7.7	3.7	2.7	6.3	2.3	4.9
EMERALD	5.3	3.3	4.7	2.7	7.0	1.3	6.3	8.3	3.7	4.0	6.7	5.0	4.9
COMPANION	7.0	7.0	4.3	4.3	4.7	4.0	5.3	7.3	3.3	2.0	5.7	3.0	4.8
GN-Z	5.7	3.7	3.0	5.3	5.3	1.0	5.7	8.3	4.3	4.3	7.7	3.7	4.8
J-37	5.7	7.0	5.3	6.0	3.7	3.3	5.7	7.0	2.7	3.7	5.3	2.7	4.8
PZA 32	5.3	7.3	3.7	6.3	4.7	2.7	5.0	7.0	2.7	2.7	6.0	3.3	4.7
CHINESE COMMON	5.3	9.0	2.7	5.3	3.3	3.0	5.7	7.3	2.0	3.0	4.7	2.3	4.5
DALZ 0101	6.0	2.3	4.0	1.0	5.0	1.0	5.7	8.3	3.7	4.3	5.7	5.0	4.3
HIMENO	5.0	9.0	1.3	8.3	4.0	1.0	4.7	3.3	2.7	2.3	4.0	4.0	4.1
DALZ 0104	4.3	1.0	.	2.7	2.0	.	2.3	3.0	3.3	1.3	4.7	3.0	2.8
6186	2.0	1.0	.	3.5	3.0	2.0	2.7	3.0	4.3	1.3	4.3	2.3	2.7
DALZ 9604	2.3	1.0	.	2.5	3.3	1.0	3.0	3.3	2.7	3.0	3.7	3.0	2.6
DALZ 0105	2.3	2.3	.	1.0	3.0	.	3.0	3.0	2.7	1.3	4.0	2.0	2.5
LSD VALUE	1.9	1.3	1.6	4.3	1.5	1.5	1.0	1.1	1.1	1.4	1.7	2.0	0.5
C.V. (%)	22.4	15.3	23.0	47.8	20.5	33.3	12.4	10.0	22.2	26.8	18.2	34.7	24.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
 STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 8B.

SPRING GREENUP RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

## SPRING GREENUP RATINGS 1-9; 9=COMPLETELY GREEN 2/

NAME	AR1	IL2	IN1	KY1	MD1	M01	MS1	NC1	NM1	OK1	SC1	TX1	MEAN
ZENITH	6.3	8.0	5.3	5.7	5.0	4.3	5.7	7.3	3.0	3.7	6.3	3.7	5.4
PZB 33	7.3	8.3	3.7	6.0	4.0	4.0	5.7	7.7	3.0	4.0	5.7	4.3	5.3
PST-R7ZM	7.3	6.3	4.3	6.0	4.0	2.3	5.7	7.7	3.3	2.3	6.7	4.0	5.0
PST-R7MA	6.3	6.3	5.0	6.3	3.3	3.7	5.0	7.7	3.7	2.7	6.3	2.3	4.9
COMPANION	7.0	7.0	4.3	4.3	4.7	4.0	5.3	7.3	3.3	2.0	5.7	3.0	4.8
J-37	5.7	7.0	5.3	6.0	3.7	3.3	5.7	7.0	2.7	3.7	5.3	2.7	4.8
PZA 32	5.3	7.3	3.7	6.3	4.7	2.7	5.0	7.0	2.7	2.7	6.0	3.3	4.7
CHINESE COMMON	5.3	9.0	2.7	5.3	3.3	3.0	5.7	7.3	2.0	3.0	4.7	2.3	4.5
LSD VALUE	1.2	1.2	1.5	5.3	1.7	1.6	0.8	1.1	1.1	1.6	1.8	2.0	0.6
C.V. (%)	11.6	9.9	22.3	57.2	25.5	28.7	9.2	9.6	22.9	32.6	19.5	38.2	26.1

TABLE 8C.

SPRING GREENUP RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

## SPRING GREENUP RATINGS 1-9; 9=COMPLETELY GREEN 2/

NAME	AR1	IL2	IN1	KY1	MD1	M01	MS1	NC1	NM1	OK1	SC1	TX1	MEAN
BMZ 230	8.0	5.3	6.3	9.0	6.0	4.0	7.0	8.0	2.7	6.0	7.3	4.7	6.2
MEYER	7.7	7.7	3.3	8.7	4.3	2.3	6.3	7.3	3.0	2.0	6.7	2.3	5.1
ZORRO	5.0	3.3	3.7	5.3	6.3	1.0	6.3	8.7	3.0	4.0	7.3	6.3	5.0
DALZ 0102	2.3	5.0	4.0	8.7	6.0	1.3	5.7	7.7	3.0	6.0	5.7	3.7	4.9
EMERALD	5.3	3.3	4.7	2.7	7.0	1.3	6.3	8.3	3.7	4.0	6.7	5.0	4.9
GN-Z	5.7	3.7	3.0	5.3	5.3	1.0	5.7	8.3	4.3	4.3	7.7	3.7	4.8
DALZ 0101	6.0	2.3	4.0	1.0	5.0	1.0	5.7	8.3	3.7	4.3	5.7	5.0	4.3
HIMENO	5.0	9.0	1.3	8.3	4.0	1.0	4.7	3.3	2.7	2.3	4.0	4.0	4.1
DALZ 0104	4.3	1.0	.	2.7	2.0	.	2.3	3.0	3.3	1.3	4.7	3.0	2.8
6186	2.0	1.0	.	3.5	3.0	2.0	2.7	3.0	4.3	1.3	4.3	2.3	2.7
DALZ 9604	2.3	1.0	.	2.5	3.3	1.0	3.0	3.3	2.7	3.0	3.7	3.0	2.6
DALZ 0105	2.3	2.3	.	1.0	3.0	.	3.0	3.0	2.7	1.3	4.0	2.0	2.5
LSD VALUE	2.3	1.3	1.6	2.9	1.4	1.1	1.1	1.0	1.1	1.2	1.6	2.0	0.5
C.V. (%)	30.5	22.2	23.7	32.9	16.8	33.4	14.5	10.3	21.8	22.9	17.2	32.7	23.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 9A.

LEAF TEXTURE RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2005 DATA

NAME	LEAF TEXTURE RATINGS 1-9; 9=VERY FINE 2/										
	AR1	IL2	IN1	MO1	MS1	NC1	NM1	OK1	SC1	TX1	MEAN
EMERALD	8.7	9.0	9.0	8.0	8.0	8.7	7.3	9.0	7.0	6.0	8.1
ZORRO	9.0	9.0	8.7	8.7	8.0	8.7	7.3	9.0	6.0	6.0	8.0
DALZ 0101	6.0	9.0	9.0	8.3	8.0	8.7	8.0	9.0	6.0	6.0	7.8
DALZ 0105	8.7	8.7	.	.	7.0	8.3	7.0	9.0	6.0	6.3	7.6
DALZ 0104	7.3	8.3	.	.	7.0	8.3	7.3	9.0	7.0	6.3	7.6
GN-Z	6.0	6.7	.	8.0	6.7	6.7	5.7	8.0	7.7	5.0	6.7
MEYER	6.7	7.7	7.0	6.3	7.0	7.0	5.0	8.7	6.7	4.7	6.7
DALZ 9604	5.3	7.0	.	.	5.7	7.0	5.7	7.7	8.0	5.0	6.4
6186	5.0	7.0	.	5.0	6.3	7.0	6.0	8.0	7.0	5.0	6.3
PST-R7ZM	5.3	6.3	6.0	6.0	5.7	5.3	4.3	8.0	9.0	4.3	6.0
PST-R7MA	5.0	6.0	6.0	6.0	5.7	6.0	4.7	7.7	9.0	4.0	6.0
DALZ 0102	7.3	4.7	4.0	5.3	6.0	4.3	3.7	6.3	8.0	4.0	5.4
ZENITH	4.7	4.7	4.3	4.7	5.7	5.7	4.0	7.0	9.0	3.7	5.3
HIMENO	4.7	4.0	3.7	6.3	5.0	5.0	4.3	7.0	9.0	4.0	5.3
PZA 32	4.0	5.0	4.3	4.7	5.0	5.3	5.0	7.0	8.3	3.7	5.2
PZB 33	4.0	4.7	3.7	5.0	5.0	5.0	4.3	7.3	9.0	4.0	5.2
CHINESE COMMON	4.0	2.0	2.7	5.0	4.7	4.3	4.3	7.0	9.0	3.7	4.7
BMZ 230	4.0	3.0	2.7	5.0	4.3	4.0	3.7	6.3	9.0	4.0	4.6
J-37	4.0	3.0	3.3	4.3	4.7	5.0	3.0	6.7	9.0	3.0	4.6
COMPANION	4.3	2.7	2.7	4.3	5.0	4.7	3.0	6.3	9.0	3.7	4.6
LSD VALUE	1.8	1.0	1.2	1.3	0.6	0.8	0.9	0.7	1.1	0.6	0.3
C.V. (%)	20.1	10.7	15.1	11.7	6.4	8.0	10.6	5.6	8.5	7.9	10.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
 STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 9B.

LEAF TEXTURE RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

LEAF TEXTURE RATINGS 1-9; 9=VERY FINE 2/											
NAME	AR1	IL2	IN1	MO1	MS1	NC1	NM1	OK1	SC1	TX1	MEAN
PST-R7ZM	5.3	6.3	6.0	6.0	5.7	5.3	4.3	8.0	9.0	4.3	6.0
PST-R7MA	5.0	6.0	6.0	6.0	5.7	6.0	4.7	7.7	9.0	4.0	6.0
ZENITH	4.7	4.7	4.3	4.7	5.7	5.7	4.0	7.0	9.0	3.7	5.3
PZA 32	4.0	5.0	4.3	4.7	5.0	5.3	5.0	7.0	8.3	3.7	5.2
PZB 33	4.0	4.7	3.7	5.0	5.0	5.0	4.3	7.3	9.0	4.0	5.2
CHINESE COMMON	4.0	2.0	2.7	5.0	4.7	4.3	4.3	7.0	9.0	3.7	4.7
J-37	4.0	3.0	3.3	4.3	4.7	5.0	3.0	6.7	9.0	3.0	4.6
COMPANION	4.3	2.7	2.7	4.3	5.0	4.7	3.0	6.3	9.0	3.7	4.6
LSD VALUE	1.1	1.3	1.5	1.1	0.7	0.9	1.0	0.7	0.3	0.7	0.3
C.V. (%)	16.0	19.0	22.7	13.3	8.8	11.2	15.8	5.7	2.3	12.2	11.9

TABLE 9C.

LEAF TEXTURE RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

LEAF TEXTURE RATINGS 1-9; 9=VERY FINE 2/											
NAME	AR1	IL2	IN1	MO1	MS1	NC1	NM1	OK1	SC1	TX1	MEAN
EMERALD	8.7	9.0	9.0	8.0	8.0	8.7	7.3	9.0	7.0	6.0	8.1
ZORRO	9.0	9.0	8.7	8.7	8.0	8.7	7.3	9.0	6.0	6.0	8.0
DALZ 0101	6.0	9.0	9.0	8.3	8.0	8.7	8.0	9.0	6.0	6.0	7.8
DALZ 0105	8.7	8.7	.	.	7.0	8.3	7.0	9.0	6.0	6.3	7.6
DALZ 0104	7.3	8.3	.	.	7.0	8.3	7.3	9.0	7.0	6.3	7.6
GN-Z	6.0	6.7	.	8.0	6.7	6.7	5.7	8.0	7.7	5.0	6.7
MEYER	6.7	7.7	7.0	6.3	7.0	7.0	5.0	8.7	6.7	4.7	6.7
DALZ 9604	5.3	7.0	.	.	5.7	7.0	5.7	7.7	8.0	5.0	6.4
6186	5.0	7.0	.	5.0	6.3	7.0	6.0	8.0	7.0	5.0	6.3
DALZ 0102	7.3	4.7	4.0	5.3	6.0	4.3	3.7	6.3	8.0	4.0	5.4
HIMENO	4.7	4.0	3.7	6.3	5.0	5.0	4.3	7.0	9.0	4.0	5.3
BMZ 230	4.0	3.0	2.7	5.0	4.3	4.0	3.7	6.3	9.0	4.0	4.6
LSD VALUE	2.2	0.8	0.9	1.4	0.5	0.7	0.8	0.7	1.4	0.5	0.4
C.V. (%)	20.8	6.7	8.5	10.7	5.1	6.3	8.0	5.5	11.7	5.6	10.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 10A. SPRING DENSITY RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2005 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	FL3	NM1	TX1	MEAN
ZORRO	8.0	9.0	8.0	8.3
DALZ 0102	8.0	9.0	7.7	8.2
DALZ 0104	8.0	9.0	7.7	8.2
DALZ 0105	8.0	9.0	7.7	8.2
GN-Z	8.0	9.0	7.7	8.2
HIMENO	8.0	9.0	7.7	8.2
BMZ 230	8.0	9.0	7.0	8.0
EMERALD	8.0	8.0	8.0	8.0
DALZ 9604	8.0	9.0	6.7	7.9
6186	7.7	9.0	6.7	7.8
DALZ 0101	8.0	7.3	7.7	7.7
CHINESE COMMON	8.0	8.7	5.7	7.4
COMPANION	8.0	9.0	5.3	7.4
ZENITH	7.7	8.3	6.0	7.3
PZA 32	6.0	9.0	6.0	7.0
PZB 33	7.0	8.3	5.3	6.9
PST-R7MA	8.0	7.3	5.3	6.9
J-37	7.0	8.3	5.0	6.8
MEYER	7.7	7.0	5.7	6.8
PST-R7ZM	8.0	7.3	4.7	6.7
LSD VALUE	1.1	0.7	1.6	0.7
C.V. (%)	9.1	4.8	15.1	9.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 10B. SPRING DENSITY RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	FL3	NM1	TX1	MEAN
CHINESE COMMON	8.0	8.7	5.7	7.4
COMPANION	8.0	9.0	5.3	7.4
ZENITH	7.7	8.3	6.0	7.3
PZA 32	6.0	9.0	6.0	7.0
PZB 33	7.0	8.3	5.3	6.9
PST-R7MA	8.0	7.3	5.3	6.9
J-37	7.0	8.3	5.0	6.8
PST-R7ZM	8.0	7.3	4.7	6.7
LSD VALUE	1.7	1.0	1.9	0.9
C.V. (%)	14.5	7.4	22.3	14.2

TABLE 10C. SPRING DENSITY RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	FL3	NM1	TX1	MEAN
ZORRO	8.0	9.0	8.0	8.3
DALZ 0102	8.0	9.0	7.7	8.2
DALZ 0104	8.0	9.0	7.7	8.2
DALZ 0105	8.0	9.0	7.7	8.2
GN-Z	8.0	9.0	7.7	8.2
HIMENO	8.0	9.0	7.7	8.2
BMZ 230	8.0	9.0	7.0	8.0
EMERALD	8.0	8.0	8.0	8.0
DALZ 9604	8.0	9.0	6.7	7.9
6186	7.7	9.0	6.7	7.8
DALZ 0101	8.0	7.3	7.7	7.7
MEYER	7.7	7.0	5.7	6.8
LSD VALUE	0.4	0.3	1.3	0.5
C.V. (%)	3.0	1.9	11.1	6.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 11A. SUMMER DENSITY RATINGS OF ZOysiAGRASS CULTIVARS 1/  
2005 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	FL3	MS1	NM1	OK1	TX1	MEAN
ZORRO	9.0	8.0	9.0	9.0	8.3	8.7
DALZ 0101	9.0	8.0	8.7	9.0	7.3	8.4
DALZ 0104	9.0	7.0	9.0	9.0	7.7	8.3
DALZ 0105	9.0	7.0	9.0	9.0	7.7	8.3
EMERALD	9.0	8.0	9.0	9.0	6.7	8.3
GN-Z	9.0	6.7	9.0	9.0	6.7	8.1
DALZ 9604	9.0	5.7	9.0	8.7	7.3	7.9
6186	9.0	6.0	9.0	7.7	7.0	7.7
DALZ 0102	9.0	6.3	9.0	7.7	6.7	7.7
HIMENO	9.0	6.0	9.0	7.3	6.0	7.5
MEYER	8.7	7.0	8.3	8.7	3.7	7.3
BMZ 230	9.0	6.0	9.0	6.3	5.7	7.2
PZB 33	9.0	5.7	9.0	7.0	5.3	7.2
PST-R7ZM	9.0	6.0	8.3	8.0	3.7	7.0
ZENITH	9.0	6.0	9.0	6.3	4.7	7.0
PST-R7MA	9.0	6.3	8.3	7.0	3.0	6.7
PZA 32	8.7	5.3	9.0	6.0	4.0	6.6
COMPANION	9.0	5.7	9.0	5.7	3.3	6.5
J-37	8.7	4.7	9.0	6.0	3.3	6.3
CHINESE COMMON	9.0	5.0	8.7	5.0	3.7	6.3
LSD VALUE	0.4	0.6	0.5	0.7	1.3	0.3
C.V. (%)	2.5	5.8	3.3	5.7	14.6	6.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 11B. SUMMER DENSITY RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	FL3	MS1	NM1	OK1	TX1	MEAN
PZB 33	9.0	5.7	9.0	7.0	5.3	7.2
PST-R7ZM	9.0	6.0	8.3	8.0	3.7	7.0
ZENITH	9.0	6.0	9.0	6.3	4.7	7.0
PST-R7MA	9.0	6.3	8.3	7.0	3.0	6.7
PZA 32	8.7	5.3	9.0	6.0	4.0	6.6
COMPANION	9.0	5.7	9.0	5.7	3.3	6.5
J-37	8.7	4.7	9.0	6.0	3.3	6.3
CHINESE COMMON	9.0	5.0	8.7	5.0	3.7	6.3
LSD VALUE	0.5	0.7	0.6	0.7	1.4	0.4
C.V. (%)	3.2	8.2	4.0	7.2	22.3	7.8

TABLE 11C. SUMMER DENSITY RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	FL3	MS1	NM1	OK1	TX1	MEAN
ZORRO	9.0	8.0	9.0	9.0	8.3	8.7
DALZ 0101	9.0	8.0	8.7	9.0	7.3	8.4
DALZ 0104	9.0	7.0	9.0	9.0	7.7	8.3
DALZ 0105	9.0	7.0	9.0	9.0	7.7	8.3
EMERALD	9.0	8.0	9.0	9.0	6.7	8.3
GN-Z	9.0	6.7	9.0	9.0	6.7	8.1
DALZ 9604	9.0	5.7	9.0	8.7	7.3	7.9
6186	9.0	6.0	9.0	7.7	7.0	7.7
DALZ 0102	9.0	6.3	9.0	7.7	6.7	7.7
HIMENO	9.0	6.0	9.0	7.3	6.0	7.5
MEYER	8.7	7.0	8.3	8.7	3.7	7.3
BMZ 230	9.0	6.0	9.0	6.3	5.7	7.2
LSD VALUE	0.3	0.5	0.4	0.7	1.3	0.3
C.V. (%)	1.9	4.2	2.6	4.9	11.6	5.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 12A. FALL DENSITY RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2005 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	FL3	NM1	TX1	MEAN
DALZ 0105	9.0	9.0	7.7	8.6
6186	9.0	9.0	7.0	8.3
DALZ 9604	9.0	9.0	7.0	8.3
DALZ 0101	9.0	9.0	6.3	8.1
DALZ 0102	9.0	9.0	6.3	8.1
DALZ 0104	9.0	9.0	6.3	8.1
EMERALD	9.0	9.0	6.3	8.1
ZORRO	8.7	9.0	6.7	8.1
GN-Z	9.0	9.0	5.7	7.9
BMZ 230	8.7	9.0	5.3	7.7
J-37	9.0	9.0	4.0	7.3
CHINESE COMMON	9.0	9.0	3.7	7.2
HIMENO	9.0	8.3	4.0	7.1
PZB 33	8.7	9.0	3.3	7.0
COMPANION	8.7	9.0	2.7	6.8
PST-R7MA	8.3	9.0	2.7	6.7
ZENITH	8.0	9.0	2.7	6.6
PZA 32	8.0	9.0	2.0	6.3
MEYER	8.0	8.0	2.3	6.1
PST-R7ZM	7.7	8.0	2.3	6.0
LSD VALUE	1.1	0.2	1.9	0.7
C.V. (%)	7.7	1.5	24.6	10.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 12B. FALL DENSITY RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	FL3	NM1	TX1	MEAN
J-37	9.0	9	4.0	7.3
CHINESE COMMON	9.0	9	3.7	7.2
PZB 33	8.7	9	3.3	7.0
COMPANION	8.7	9	2.7	6.8
PST-R7MA	8.3	9	2.7	6.7
ZENITH	8.0	9	2.7	6.6
PZA 32	8.0	9	2.0	6.3
PST-R7ZM	7.7	8	2.3	6.0
LSD VALUE	1.3	0	1.9	0.8
C.V. (%)	9.7	0	40.2	12.2

TABLE 12C. FALL DENSITY RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	FL3	NM1	TX1	MEAN
DALZ 0105	9.0	9.0	7.7	8.6
6186	9.0	9.0	7.0	8.3
DALZ 9604	9.0	9.0	7.0	8.3
DALZ 0101	9.0	9.0	6.3	8.1
DALZ 0102	9.0	9.0	6.3	8.1
DALZ 0104	9.0	9.0	6.3	8.1
EMERALD	9.0	9.0	6.3	8.1
ZORRO	8.7	9.0	6.7	8.1
GN-Z	9.0	9.0	5.7	7.9
BMZ 230	8.7	9.0	5.3	7.7
HIMENO	9.0	8.3	4.0	7.1
MEYER	8.0	8.0	2.3	6.1
LSD VALUE	0.9	0.3	1.9	0.7
C.V. (%)	6.2	1.9	19.5	9.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 13A. PERCENT LIVING GROUND COVER (SPRING) RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2005 DATA

PERCENT LIVING GROUND COVER IN SPRING: LOCATIONS 2/

NAME	IL2
CHINESE COMMON	99.0
BMZ 230	98.3
COMPANION	97.7
DALZ 0102	97.7
HIMENO	97.7
J-37	97.7
ZENITH	97.7
PZB 33	96.7
PZA 32	94.7
ZORRO	93.7
MEYER	92.7
PST-R7MA	92.0
GN-Z	84.3
EMERALD	82.7
PST-R7ZM	73.0
DALZ 0101	64.7
DALZ 0104	45.3
DALZ 0105	20.0
DALZ 9604	11.0
6186	4.0
LSD VALUE	15.8
C.V. (%)	12.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 13B. PERCENT LIVING GROUND COVER (SPRING) RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

PERCENT LIVING GROUND COVER IN SPRING: LOCATIONS 2/

NAME	IL2
CHINESE COMMON	99.0
COMPANION	97.7
J-37	97.7
ZENITH	97.7
PZB 33	96.7
PZA 32	94.7
PST-R7MA	92.0
PST-R7ZM	73.0
LSD VALUE	5.6
C.V. (%)	3.7

TABLE 13C. PERCENT LIVING GROUND COVER (SPRING) RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

PERCENT LIVING GROUND COVER IN SPRING: LOCATIONS 2/

NAME	IL2
BMZ 230	98.3
DALZ 0102	97.7
HIMENO	97.7
ZORRO	93.7
MEYER	92.7
GN-Z	84.3
EMERALD	82.7
DALZ 0101	64.7
DALZ 0104	45.3
DALZ 0105	20.0
DALZ 9604	11.0
6186	4.0
LSD VALUE	19.8
C.V. (%)	18.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 14A. PERCENT LIVING GROUND COVER (SUMMER) RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2005 DATA

PERCENT LIVING GROUND COVER IN SUMMER: LOCATIONS 2/

NAME	IL2	OK1	MEAN
DALZ 0102	99.0	99.0	99.0
EMERALD	99.0	98.3	98.7
ZORRO	98.3	99.0	98.7
GN-Z	99.0	97.0	98.0
MEYER	98.3	97.0	97.7
DALZ 0101	94.7	99.0	96.8
HIMENO	99.0	93.3	96.2
J-37	97.7	93.0	95.3
BMZ 230	99.0	90.3	94.7
PST-R7MA	93.3	94.0	93.7
PZA 32	95.3	90.7	93.0
PZB 33	99.0	87.0	93.0
DALZ 0104	86.7	98.7	92.7
COMPANION	93.0	87.7	90.3
ZENITH	95.7	83.7	89.7
CHINESE COMMON	99.0	63.3	81.2
DALZ 0105	63.3	99.0	81.2
PST-R7ZM	61.3	86.7	74.0
DALZ 9604	24.0	98.0	61.0
6186	8.3	98.3	53.3
LSD VALUE	18.7	13.5	11.5
C.V. (%)	13.7	9.0	11.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 14B. PERCENT LIVING GROUND COVER (SUMMER) RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

PERCENT LIVING GROUND COVER IN SUMMER: LOCATIONS 2/

NAME	IL2	OK1	MEAN
J-37	97.7	93.0	95.3
PST-R7MA	93.3	94.0	93.7
PZA 32	95.3	90.7	93.0
PZB 33	99.0	87.0	93.0
COMPANION	93.0	87.7	90.3
ZENITH	95.7	83.7	89.7
CHINESE COMMON	99.0	63.3	81.2
PST-R7ZM	61.3	86.7	74.0
LSD VALUE	13.7	19.8	12.0
C.V. (%)	9.3	14.4	11.9

TABLE 14C. PERCENT LIVING GROUND COVER (SUMMER) RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

PERCENT LIVING GROUND COVER IN SUMMER: LOCATIONS 2/

NAME	IL2	OK1	MEAN
DALZ 0102	99.0	99.0	99.0
EMERALD	99.0	98.3	98.7
ZORRO	98.3	99.0	98.7
GN-Z	99.0	97.0	98.0
MEYER	98.3	97.0	97.7
DALZ 0101	94.7	99.0	96.8
HIMENO	99.0	93.3	96.2
BMZ 230	99.0	90.3	94.7
DALZ 0104	86.7	98.7	92.7
DALZ 0105	63.3	99.0	81.2
DALZ 9604	24.0	98.0	61.0
6186	8.3	98.3	53.3
LSD VALUE	21.4	6.4	11.2
C.V. (%)	16.5	4.1	11.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 15A. PERCENT LIVING GROUND COVER (FALL) RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2005 DATA

PERCENT LIVING GROUND COVER IN FALL: LOCATIONS 2/

NAME	IL2	OK1	MEAN
ZORRO	99.0	99.0	99.0
BMZ 230	99.0	98.7	98.8
DALZ 0102	99.0	98.3	98.7
EMERALD	99.0	98.3	98.7
GN-Z	99.0	98.3	98.7
MEYER	99.0	98.3	98.7
PST-R7MA	99.0	98.3	98.7
DALZ 0104	97.0	99.0	98.0
DALZ 0101	96.0	99.0	97.5
HIMENO	99.0	93.7	96.3
PZA 32	94.7	97.0	95.8
COMPANION	99.0	89.7	94.3
PST-R7ZM	91.0	95.0	93.0
ZENITH	99.0	85.0	92.0
PZB 33	99.0	84.0	91.5
CHINESE COMMON	99.0	77.7	88.3
DALZ 0105	74.0	99.0	86.5
DALZ 9604	70.3	96.7	83.5
6186	30.7	98.3	64.5
J-37	99.0	28.3	63.7
LSD VALUE	18.5	13.2	11.4
C.V. (%)	12.5	9.0	10.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 15B. PERCENT LIVING GROUND COVER (FALL) RATINGS OF ZOysiagrass (SEEDED) CULTIVARS 1/  
2005 DATA

PERCENT LIVING GROUND COVER IN FALL: LOCATIONS 2/

NAME	IL2	OK1	MEAN
PST-R7MA	99.0	98.3	98.7
PZA 32	94.7	97.0	95.8
COMPANION	99.0	89.7	94.3
PST-R7ZM	91.0	95.0	93.0
ZENITH	99.0	85.0	92.0
PZB 33	99.0	84.0	91.5
CHINESE COMMON	99.0	77.7	88.3
J-37	99.0	28.3	63.7
LSD VALUE	6.2	20.7	10.8
C.V. (%)	3.9	15.7	10.6

TABLE 15C. PERCENT LIVING GROUND COVER (FALL) RATINGS OF ZOysiagrass (VEGETATIVE) CULTIVARS 1/  
2005 DATA

PERCENT LIVING GROUND COVER IN FALL: LOCATIONS 2/

NAME	IL2	OK1	MEAN
ZORRO	99.0	99.0	99.0
BMZ 230	99.0	98.7	98.8
DALZ 0102	99.0	98.3	98.7
EMERALD	99.0	98.3	98.7
GN-Z	99.0	98.3	98.7
MEYER	99.0	98.3	98.7
DALZ 0104	97.0	99.0	98.0
DALZ 0101	96.0	99.0	97.5
HIMENO	99.0	93.7	96.3
DALZ 0105	74.0	99.0	86.5
DALZ 9604	70.3	96.7	83.5
6186	30.7	98.3	64.5
LSD VALUE	23.3	2.7	11.7
C.V. (%)	16.4	1.7	11.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 16A. FROST TOLERANCE RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2005 DATA

FROST TOLERANCE RATINGS 1-9; 9=NO INJURY 2/

NAME	AR1	IL2	KY1	M01	OK1	MEAN
PZA 32	5.0	7.7	9.0	5.3	9.0	7.2
PZB 33	4.7	7.0	9.0	5.7	9.0	7.1
ZENITH	4.3	7.0	9.0	5.0	9.0	6.9
PST-R7MA	4.3	6.7	8.0	5.7	9.0	6.7
COMPANION	5.3	7.3	8.7	4.7	7.7	6.7
MEYER	6.3	4.3	9.0	5.0	9.0	6.7
6186	4.0	9.0	4.5	7.0	9.0	6.7
ZORRO	7.3	6.0	9.0	1.7	9.0	6.6
BMZ 230	5.7	7.0	8.0	2.7	9.0	6.5
DALZ 0102	6.0	6.7	7.7	2.3	9.0	6.3
HIMENO	6.3	6.0	9.0	4.0	6.3	6.3
PST-R7ZM	4.3	5.0	8.3	5.0	9.0	6.3
DALZ 0105	4.7	7.0	4.5	.	9.0	6.3
GN-Z	4.3	6.7	9.0	2.0	9.0	6.2
EMERALD	7.3	5.0	6.3	3.0	9.0	6.1
DALZ 9604	7.0	4.7	5.0	.	7.7	6.1
DALZ 0101	4.7	5.3	7.7	2.3	9.0	5.8
J-37	4.0	5.7	8.7	5.0	4.0	5.5
DALZ 0104	6.3	2.3	4.0	.	9.0	5.4
CHINESE COMMON	3.7	3.3	8.7	5.0	5.0	5.1
LSD VALUE	1.6	1.3	1.4	2.1	0.6	0.6
C.V. (%)	19.4	13.8	10.9	27.0	4.7	13.6

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 16B. FROST TOLERANCE RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

FROST TOLERANCE RATINGS 1-9; 9=NO INJURY 2/

NAME	AR1	IL2	KY1	M01	OK1	MEAN
PZA 32	5.0	7.7	9.0	5.3	9.0	7.2
PZB 33	4.7	7.0	9.0	5.7	9.0	7.1
ZENITH	4.3	7.0	9.0	5.0	9.0	6.9
PST-R7MA	4.3	6.7	8.0	5.7	9.0	6.7
COMPANION	5.3	7.3	8.7	4.7	7.7	6.7
PST-R7ZM	4.3	5.0	8.3	5.0	9.0	6.3
J-37	4.0	5.7	8.7	5.0	4.0	5.5
CHINESE COMMON	3.7	3.3	8.7	5.0	5.0	5.1
LSD VALUE	1.0	1.4	0.7	2.0	0.3	0.5
C.V. (%)	13.7	13.6	4.7	23.8	2.6	11.5

TABLE 16C. FROST TOLERANCE RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

FROST TOLERANCE RATINGS 1-9; 9=NO INJURY 2/

NAME	AR1	IL2	KY1	M01	OK1	MEAN
MEYER	6.3	4.3	9.0	5.0	9.0	6.7
6186	4.0	9.0	4.5	7.0	9.0	6.7
ZORRO	7.3	6.0	9.0	1.7	9.0	6.6
BMZ 230	5.7	7.0	8.0	2.7	9.0	6.5
DALZ 0102	6.0	6.7	7.7	2.3	9.0	6.3
HIMENO	6.3	6.0	9.0	4.0	6.3	6.3
DALZ 0105	4.7	7.0	4.5	.	9.0	6.3
GN-Z	4.3	6.7	9.0	2.0	9.0	6.2
EMERALD	7.3	5.0	6.3	3.0	9.0	6.1
DALZ 9604	7.0	4.7	5.0	.	7.7	6.1
DALZ 0101	4.7	5.3	7.7	2.3	9.0	5.8
DALZ 0104	6.3	2.3	4.0	.	9.0	5.4
LSD VALUE	2.0	1.3	1.8	2.0	0.8	0.7
C.V. (%)	21.0	14.0	15.0	30.7	5.4	15.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 17A. WINTER COLOR RATINGS OF ZOysiAGRASS CULTIVARS 1/  
2005 DATA

WINTER COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	CA3	FL3	NM1	OK1	MEAN
6186	5.3	2.3	1.0	8.0	4.2
DALZ 9604	5.3	2.3	1.0	8.0	4.2
DALZ 0104	4.7	2.3	1.0	8.0	4.0
DALZ 0105	5.0	2.0	1.0	7.0	3.8
HIMENO	.	1.0	1.0	9.0	3.7
DALZ 0101	4.7	1.0	1.3	6.3	3.3
DALZ 0102	1.0	1.0	1.0	9.0	3.0
EMERALD	2.3	1.3	1.3	7.0	3.0
ZENITH	1.0	1.0	1.3	8.7	3.0
CHINESE COMMON	1.0	1.0	1.0	8.0	2.8
PST-R7MA	1.0	1.0	1.0	8.0	2.8
PZB 33	1.0	1.0	1.0	8.0	2.8
ZORRO	1.7	1.0	1.0	6.7	2.6
GN-Z	1.0	1.3	1.0	7.0	2.6
COMPANION	1.0	1.0	1.0	7.0	2.5
PZA 32	1.0	1.0	1.0	7.0	2.5
MEYER	1.0	1.0	1.0	6.3	2.3
J-37	1.0	1.0	1.0	6.0	2.3
PST-R7ZM	1.0	1.0	1.0	6.0	2.3
BMZ 230	1.0	1.0	1.0	5.3	2.1
LSD VALUE	0.6	0.5	0.4	0.6	0.3
C.V. (%)	15.9	22.5	21.3	5.0	10.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 17B. WINTER COLOR RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

WINTER COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	CA3	FL3	NM1	OK1	MEAN
ZENITH	1	1	1.3	8.7	3.0
CHINESE COMMON	1	1	1.0	8.0	2.8
PST-R7MA	1	1	1.0	8.0	2.8
PZB 33	1	1	1.0	8.0	2.8
COMPANION	1	1	1.0	7.0	2.5
PZA 32	1	1	1.0	7.0	2.5
J-37	1	1	1.0	6.0	2.3
PST-R7ZM	1	1	1.0	6.0	2.3
LSD VALUE	0	0	0.3	0.7	0.2
C.V. (%)	0	0	19.6	5.6	9.1

TABLE 17C. WINTER COLOR RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

WINTER COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	CA3	FL3	NM1	OK1	MEAN
6186	5.3	2.3	1.0	8.0	4.2
DALZ 9604	5.3	2.3	1.0	8.0	4.2
DALZ 0104	4.7	2.3	1.0	8.0	4.0
DALZ 0105	5.0	2.0	1.0	7.0	3.8
HIMENO	.	1.0	1.0	9.0	3.7
DALZ 0101	4.7	1.0	1.3	6.3	3.3
DALZ 0102	1.0	1.0	1.0	9.0	3.0
EMERALD	2.3	1.3	1.3	7.0	3.0
ZORRO	1.7	1.0	1.0	6.7	2.6
GN-Z	1.0	1.3	1.0	7.0	2.6
MEYER	1.0	1.0	1.0	6.3	2.3
BMZ 230	1.0	1.0	1.0	5.3	2.1
LSD VALUE	0.7	0.6	0.4	0.5	0.3
C.V. (%)	14.2	25.3	22.3	4.6	10.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 18A. DOLLAR SPOT RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2005 DATA

DOLLAR SPOT RATINGS 1-9; 9=NO DISEASE 2/

NAME	SC1
COMPANION	9.0
DALZ 0101	9.0
EMERALD	9.0
HIMENO	9.0
PST-R7ZM	8.7
DALZ 0102	8.3
J-37	8.3
ZENITH	8.3
BMZ 230	8.0
CHINESE COMMON	8.0
DALZ 0104	8.0
GN-Z	8.0
PZA 32	8.0
ZORRO	8.0
MEYER	7.7
PZB 33	7.7
6186	7.3
DALZ 0105	7.3
DALZ 9604	7.3
PST-R7MA	6.7
LSD VALUE	1.9
C.V. (%)	14.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 18B. DOLLAR SPOT RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

DOLLAR SPOT RATINGS 1-9; 9=NO DISEASE 2/

NAME	SC1
COMPANION	9.0
PST-R7ZM	8.7
J-37	8.3
ZENITH	8.3
CHINESE COMMON	8.0
PZA 32	8.0
PZB 33	7.7
PST-R7MA	6.7
LSD VALUE	1.9
C.V. (%)	14.3

TABLE 18C. DOLLAR SPOT RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

DOLLAR SPOT RATINGS 1-9; 9=NO DISEASE 2/

NAME	SC1
DALZ 0101	9.0
EMERALD	9.0
HIMENO	9.0
DALZ 0102	8.3
BMZ 230	8.0
DALZ 0104	8.0
GN-Z	8.0
ZORRO	8.0
MEYER	7.7
6186	7.3
DALZ 0105	7.3
DALZ 9604	7.3
LSD VALUE	2.0
C.V. (%)	15.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 19A. BROWN PATCH (WARM TEMPERATURE) RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2005 DATA

BROWN PATCH RATINGS 1-9; 9=NO DISEASE 2/

NAME	SC1
ZORRO	8.0
PST-R7ZM	7.7
DALZ 0102	7.3
MEYER	7.3
ZENITH	7.3
EMERALD	7.0
GN-Z	7.0
HIMENO	7.0
BMZ 230	6.7
DALZ 0105	6.7
PST-R7MA	6.7
DALZ 0101	6.3
PZA 32	6.3
DALZ 0104	6.0
COMPANION	5.7
DALZ 9604	5.7
J-37	5.7
PZB 33	5.7
6186	4.7
CHINESE COMMON	4.7
LSD VALUE	2.2
C.V. (%)	21.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 19B. BROWN PATCH (WARM TEMPERATURE) RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

BROWN PATCH RATINGS 1-9; 9=NO DISEASE 2/

NAME	SC1
PST-R7ZM	7.7
ZENITH	7.3
PST-R7MA	6.7
PZA 32	6.3
COMPANION	5.7
J-37	5.7
PZB 33	5.7
CHINESE COMMON	4.7
LSD VALUE	3.0
C.V. (%)	30.0

TABLE 19C. BROWN PATCH (WARM TEMPERATURE) RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

BROWN PATCH RATINGS 1-9; 9=NO DISEASE 2/

NAME	SC1
ZORRO	8.0
DALZ 0102	7.3
MEYER	7.3
EMERALD	7.0
GN-Z	7.0
HIMENO	7.0
BMZ 230	6.7
DALZ 0105	6.7
DALZ 0101	6.3
DALZ 0104	6.0
DALZ 9604	5.7
6186	4.7
LSD VALUE	1.5
C.V. (%)	14.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 20A. TAKE-ALL PATCH RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2005 DATA

TAKE-ALL PATCH RATINGS 1-9; 9=NO DISEASE 2/

NAME	TX1
ZORRO	9.0
6186	8.7
DALZ 0101	8.7
DALZ 0102	8.7
DALZ 0104	8.7
DALZ 0105	8.7
DALZ 9604	8.7
EMERALD	8.7
GN-Z	8.7
BMZ 230	8.3
CHINESE COMMON	5.7
COMPANION	5.7
HIMENO	5.0
J-37	4.7
PST-R7ZM	4.3
PZA 32	4.0
PZB 33	4.0
MEYER	3.7
PST-R7MA	3.3
ZENITH	2.7
LSD VALUE	3.0
C.V. (%)	28.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 20B. TAKE-ALL PATCH RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

TAKE-ALL PATCH RATINGS 1-9; 9=NO DISEASE 2/

NAME	TX1
CHINESE COMMON	5.7
COMPANION	5.7
J-37	4.7
PST-R7ZM	4.3
PZA 32	4.0
PZB 33	4.0
PST-R7MA	3.3
ZENITH	2.7
LSD VALUE	3.9
C.V. (%)	56.5

TABLE 20C. TAKE-ALL PATCH RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

TAKE-ALL PATCH RATINGS 1-9; 9=NO DISEASE 2/

NAME	TX1
ZORRO	9.0
6186	8.7
DALZ 0101	8.7
DALZ 0102	8.7
DALZ 0104	8.7
DALZ 0105	8.7
DALZ 9604	8.7
EMERALD	8.7
GN-Z	8.7
BMZ 230	8.3
HIMENO	5.0
MEYER	3.7
LSD VALUE	2.1
C.V. (%)	16.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 21A. MOLE CRICKET RATINGS OF ZOysiAGRASS CULTIVARS 1/  
2005 DATA

MOLE CRICKET RATINGS 1-9; 9=NO DAMAGE 2/

NAME	SC1
DALZ 0102	8.3
J-37	8.3
ZENITH	8.3
DALZ 0104	8.0
DALZ 9604	8.0
ZORRO	8.0
DALZ 0105	7.7
EMERALD	7.7
MEYER	7.7
6186	7.3
BMZ 230	7.3
COMPANION	7.3
HIMENO	7.3
PST-R7ZM	7.3
GN-Z	7.0
PZB 33	7.0
PST-R7MA	6.3
PZA 32	6.3
DALZ 0101	6.0
CHINESE COMMON	5.7
LSD VALUE	2.0
C.V. (%)	16.6

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 21B. MOLE CRICKET RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

MOLE CRICKET RATINGS 1-9; 9=NO DAMAGE 2/

NAME	SC1
J-37	8.3
ZENITH	8.3
COMPANION	7.3
PST-R7ZM	7.3
PZB 33	7.0
PST-R7MA	6.3
PZA 32	6.3
CHINESE COMMON	5.7
LSD VALUE	1.9
C.V. (%)	16.8

TABLE 21C. MOLE CRICKET RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

MOLE CRICKET RATINGS 1-9; 9=NO DAMAGE 2/

NAME	SC1
DALZ 0102	8.3
DALZ 0104	8.0
DALZ 9604	8.0
ZORRO	8.0
DALZ 0105	7.7
EMERALD	7.7
MEYER	7.7
6186	7.3
BMZ 230	7.3
HIMENO	7.3
GN-Z	7.0
DALZ 0101	6.0
LSD VALUE	2.0
C.V. (%)	16.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 22A. FALL COLOR (SEPTEMBER) RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2005 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	FL3	GA1	NM1	TX1	MEAN
DALZ 0101	7.0	8.7	5.7	7.0	7.1
EMERALD	6.7	8.0	5.3	8.0	7.0
6186	6.7	8.0	5.3	7.3	6.8
HIMENO	6.7	8.0	5.7	6.0	6.6
DALZ 0105	6.3	7.7	5.0	7.3	6.6
ZORRO	7.3	7.7	5.3	6.0	6.6
GN-Z	6.7	6.7	5.3	7.0	6.4
BMZ 230	7.0	6.3	5.0	7.3	6.4
DALZ 0104	6.3	8.0	5.3	6.0	6.4
DALZ 9604	6.3	7.3	5.0	7.0	6.4
DALZ 0102	6.0	6.7	5.0	7.0	6.2
J-37	7.3	6.0	5.3	5.7	6.1
MEYER	7.0	7.3	5.3	4.7	6.1
COMPANION	6.7	7.0	5.0	5.3	6.0
PZA 32	7.0	6.7	5.3	5.0	6.0
CHINESE COMMON	6.0	5.7	5.0	6.3	5.8
PST-R7MA	6.7	6.7	4.7	5.0	5.8
PZB 33	7.3	5.7	5.7	4.3	5.8
PST-R7ZM	6.7	6.3	5.0	4.7	5.7
ZENITH	6.7	6.0	5.3	4.0	5.5
LSD VALUE	1.1	1.0	1.0	1.9	0.7
C.V. (%)	9.8	9.2	12.3	19.2	12.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 22B. FALL COLOR (SEPTEMBER) RATINGS OF ZOysiAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	FL3	GA1	NM1	TX1	MEAN
J-37	7.3	6.0	5.3	5.7	6.1
COMPANION	6.7	7.0	5.0	5.3	6.0
PZA 32	7.0	6.7	5.3	5.0	6.0
CHINESE COMMON	6.0	5.7	5.0	6.3	5.8
PST-R7MA	6.7	6.7	4.7	5.0	5.8
PZB 33	7.3	5.7	5.7	4.3	5.8
PST-R7ZM	6.7	6.3	5.0	4.7	5.7
ZENITH	6.7	6.0	5.3	4.0	5.5
LSD VALUE	1.0	1.2	0.7	2.3	0.7
C.V. (%)	9.0	12.2	8.8	28.6	15.5

TABLE 22C. FALL COLOR (SEPTEMBER) RATINGS OF ZOysiAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	FL3	GA1	NM1	TX1	MEAN
DALZ 0101	7.0	8.7	5.7	7.0	7.1
EMERALD	6.7	8.0	5.3	8.0	7.0
6186	6.7	8.0	5.3	7.3	6.8
HIMENO	6.7	8.0	5.7	6.0	6.6
DALZ 0105	6.3	7.7	5.0	7.3	6.6
ZORRO	7.3	7.7	5.3	6.0	6.6
GN-Z	6.7	6.7	5.3	7.0	6.4
BMZ 230	7.0	6.3	5.0	7.3	6.4
DALZ 0104	6.3	8.0	5.3	6.0	6.4
DALZ 9604	6.3	7.3	5.0	7.0	6.4
DALZ 0102	6.0	6.7	5.0	7.0	6.2
MEYER	7.0	7.3	5.3	4.7	6.1
LSD VALUE	1.1	0.9	1.2	1.5	0.6
C.V. (%)	10.3	7.3	14.1	13.8	11.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 23A. FALL COLOR (OCTOBER) RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2005 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	FL3	GA1	KS1	M01	NM1	SC1	TX1	MEAN
DALZ 0101	7.7	8.0	7.0	8.0	5.3	8.0	6.0	7.1
EMERALD	7.0	7.7	7.0	5.3	5.0	8.7	7.0	6.8
BMZ 230	7.3	6.7	4.0	6.0	5.0	8.7	8.0	6.5
ZORRO	7.0	7.7	5.3	6.7	5.0	8.3	5.7	6.5
DALZ 0105	7.7	7.7	2.3	.	5.0	8.0	7.3	6.3
DALZ 0104	7.3	7.7	1.7	.	5.3	8.7	6.7	6.2
HIMENO	7.0	7.7	3.3	5.7	5.3	8.7	5.7	6.2
DALZ 0102	6.7	6.0	6.0	4.0	4.3	8.7	7.3	6.1
6186	7.0	8.0	1.7	4.0	5.3	8.3	7.7	6.0
PZB 33	6.7	5.7	4.0	6.3	5.7	8.3	5.0	6.0
GN-Z	7.3	6.0	3.0	5.0	5.3	8.0	7.0	6.0
DALZ 9604	6.7	6.7	2.0	.	5.0	8.3	6.7	5.9
J-37	6.7	5.3	3.3	5.7	5.3	7.3	6.0	5.7
COMPANION	5.7	6.0	2.3	5.7	5.3	8.3	6.0	5.6
PZA 32	6.3	5.7	2.7	6.0	5.3	8.3	5.0	5.6
ZENITH	6.7	6.3	3.3	4.7	5.3	7.7	4.3	5.5
PST-R7MA	6.3	5.7	2.7	4.7	4.7	8.7	5.3	5.4
CHINESE COMMON	6.0	5.7	3.3	4.7	4.7	6.7	6.3	5.3
PST-R7ZM	5.7	5.7	2.0	5.0	4.7	8.0	6.0	5.3
MEYER	6.7	6.3	2.0	3.3	5.3	8.3	4.7	5.2
LSD VALUE	1.1	1.0	1.4	2.4	1.1	1.3	1.7	0.5
C.V. (%)	10.4	9.0	25.7	25.3	13.1	9.6	16.8	14.6

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 23B. FALL COLOR (OCTOBER) RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

NAME	FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/							
	FL3	GA1	KS1	M01	NM1	SC1	TX1	MEAN
PZB 33	6.7	5.7	4.0	6.3	5.7	8.3	5.0	6.0
J-37	6.7	5.3	3.3	5.7	5.3	7.3	6.0	5.7
COMPANION	5.7	6.0	2.3	5.7	5.3	8.3	6.0	5.6
PZA 32	6.3	5.7	2.7	6.0	5.3	8.3	5.0	5.6
ZENITH	6.7	6.3	3.3	4.7	5.3	7.7	4.3	5.5
PST-R7MA	6.3	5.7	2.7	4.7	4.7	8.7	5.3	5.4
CHINESE COMMON	6.0	5.7	3.3	4.7	4.7	6.7	6.3	5.3
PST-R7ZM	5.7	5.7	2.0	5.0	4.7	8.0	6.0	5.3
LSD VALUE	1.2	0.9	1.4	2.2	0.9	1.2	1.9	0.5
C.V. (%)	11.8	9.4	29.3	24.7	11.3	9.3	21.3	16.0

TABLE 23C. FALL COLOR (OCTOBER) RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

NAME	FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/							
	FL3	GA1	KS1	M01	NM1	SC1	TX1	MEAN
DALZ 0101	7.7	8.0	7.0	8.0	5.3	8.0	6.0	7.1
EMERALD	7.0	7.7	7.0	5.3	5.0	8.7	7.0	6.8
BMZ 230	7.3	6.7	4.0	6.0	5.0	8.7	8.0	6.5
ZORRO	7.0	7.7	5.3	6.7	5.0	8.3	5.7	6.5
DALZ 0105	7.7	7.7	2.3	.	5.0	8.0	7.3	6.3
DALZ 0104	7.3	7.7	1.7	.	5.3	8.7	6.7	6.2
HIMENO	7.0	7.7	3.3	5.7	5.3	8.7	5.7	6.2
DALZ 0102	6.7	6.0	6.0	4.0	4.3	8.7	7.3	6.1
6186	7.0	8.0	1.7	4.0	5.3	8.3	7.7	6.0
GN-Z	7.3	6.0	3.0	5.0	5.3	8.0	7.0	6.0
DALZ 9604	6.7	6.7	2.0	.	5.0	8.3	6.7	5.9
MEYER	6.7	6.3	2.0	3.3	5.3	8.3	4.7	5.2
LSD VALUE	1.1	1.0	1.4	2.7	1.2	1.3	1.5	0.5
C.V. (%)	9.7	8.7	23.8	25.9	14.2	9.7	14.2	13.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 24A. FALL COLOR (NOVEMBER) RATINGS OF ZOysiagrass CULTIVARS 1/  
2005 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	FL3	MS1	NC1	NM1	OK1	TX1	MEAN
DALZ 0101	4.0	4.0	6.3	5.7	8.3	5.3	5.6
6186	4.3	5.0	6.3	5.3	3.0	7.7	5.3
GN-Z	3.7	3.3	6.3	5.0	6.3	6.3	5.2
EMERALD	4.0	3.7	6.7	5.0	4.3	5.7	4.9
ZORRO	4.0	4.0	6.3	4.0	4.0	6.0	4.7
DALZ 0104	4.0	4.0	6.3	4.7	2.0	6.0	4.5
DALZ 0105	4.0	3.3	7.0	4.7	1.7	6.3	4.5
DALZ 9604	4.0	3.0	6.0	4.7	2.0	6.0	4.3
BMZ 230	3.7	3.3	6.7	4.0	2.0	3.0	3.8
DALZ 0102	2.7	3.0	5.0	4.7	2.7	4.0	3.7
HIMENO	3.0	3.0	5.3	4.7	1.7	3.0	3.4
J-37	5.0	2.3	4.0	5.0	1.0	2.7	3.3
MEYER	3.7	3.0	5.0	4.7	1.3	2.0	3.3
ZENITH	3.7	3.0	5.0	5.0	1.0	2.0	3.3
PZB 33	2.7	2.7	5.0	5.7	1.0	2.3	3.2
PZA 32	2.7	3.0	5.0	5.3	1.0	2.0	3.2
PST-R7MA	3.0	2.7	4.3	4.3	1.0	2.3	2.9
COMPANION	2.7	2.3	4.3	4.7	1.0	2.3	2.9
CHINESE COMMON	2.0	3.0	4.0	4.0	1.0	2.7	2.8
PST-R7ZM	2.7	2.3	4.0	4.3	1.0	2.0	2.7
LSD VALUE	0.7	0.6	1.0	1.2	1.3	1.3	0.4
C.V. (%)	13.4	12.1	11.1	15.8	33.2	19.7	16.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 24B. FALL COLOR (NOVEMBER) RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	FL3	MS1	NC1	NM1	OK1	TX1	MEAN
J-37	5.0	2.3	4.0	5.0	1	2.7	3.3
ZENITH	3.7	3.0	5.0	5.0	1	2.0	3.3
PZB 33	2.7	2.7	5.0	5.7	1	2.3	3.2
PZA 32	2.7	3.0	5.0	5.3	1	2.0	3.2
PST-R7MA	3.0	2.7	4.3	4.3	1	2.3	2.9
COMPANION	2.7	2.3	4.3	4.7	1	2.3	2.9
CHINESE COMMON	2.0	3.0	4.0	4.0	1	2.7	2.8
PST-R7ZM	2.7	2.3	4.0	4.3	1	2.0	2.7
LSD VALUE	0.9	0.7	0.7	1.2	0	0.7	0.3
C.V. (%)	19.0	17.1	10.2	15.9	0	19.9	16.7

TABLE 24C. FALL COLOR (NOVEMBER) RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	FL3	MS1	NC1	NM1	OK1	TX1	MEAN
DALZ 0101	4.0	4.0	6.3	5.7	8.3	5.3	5.6
6186	4.3	5.0	6.3	5.3	3.0	7.7	5.3
GN-Z	3.7	3.3	6.3	5.0	6.3	6.3	5.2
EMERALD	4.0	3.7	6.7	5.0	4.3	5.7	4.9
ZORRO	4.0	4.0	6.3	4.0	4.0	6.0	4.7
DALZ 0104	4.0	4.0	6.3	4.7	2.0	6.0	4.5
DALZ 0105	4.0	3.3	7.0	4.7	1.7	6.3	4.5
DALZ 9604	4.0	3.0	6.0	4.7	2.0	6.0	4.3
BMZ 230	3.7	3.3	6.7	4.0	2.0	3.0	3.8
DALZ 0102	2.7	3.0	5.0	4.7	2.7	4.0	3.7
HIMENO	3.0	3.0	5.3	4.7	1.7	3.0	3.4
MEYER	3.7	3.0	5.0	4.7	1.3	2.0	3.3
LSD VALUE	0.6	0.5	1.1	1.2	1.6	1.5	0.5
C.V. (%)	9.9	9.4	11.2	15.7	30.9	18.4	16.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 25A. FALL COLOR (DECEMBER) RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2005 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	CA3	NM1	TX1	MEAN
DALZ 0104	4.7	4.0	2.7	3.8
GN-Z	4.0	4.3	3.0	3.8
ZORRO	3.7	4.0	2.7	3.4
DALZ 0101	2.7	4.3	3.0	3.3
DALZ 9604	4.0	3.3	2.3	3.2
6186	4.3	4.3	1.0	3.2
HIMENO	.	4.0	2.3	3.2
EMERALD	2.3	4.0	3.0	3.1
DALZ 0105	3.0	3.7	2.3	3.0
DALZ 0102	2.3	3.7	2.7	2.9
BMZ 230	1.3	4.0	2.0	2.4
J-37	1.0	4.3	2.0	2.4
PZA 32	1.0	4.3	1.7	2.3
ZENITH	1.0	4.0	1.7	2.2
CHINESE COMMON	1.0	3.7	2.0	2.2
MEYER	1.0	4.3	1.3	2.2
PZB 33	1.0	4.3	1.3	2.2
PST-R7ZM	1.0	3.7	1.7	2.1
COMPANION	1.0	3.7	1.3	2.0
PST-R7MA	1.0	3.7	1.3	2.0
LSD VALUE	1.1	1.1	0.9	0.6
C.V. (%)	29.8	16.5	27.2	22.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 25B. FALL COLOR (DECEMBER) RATINGS OF ZOysiAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	CA3	NM1	TX1	MEAN
J-37	1	4.3	2.0	2.4
PZA 32	1	4.3	1.7	2.3
ZENITH	1	4.0	1.7	2.2
CHINESE COMMON	1	3.7	2.0	2.2
PZB 33	1	4.3	1.3	2.2
PST-R7ZM	1	3.7	1.7	2.1
COMPANION	1	3.7	1.3	2.0
PST-R7MA	1	3.7	1.3	2.0
LSD VALUE	0	1.0	0.8	0.5
C.V. (%)	0	16.3	30.8	22.2

TABLE 25C. FALL COLOR (DECEMBER) RATINGS OF ZOysiAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	CA3	NM1	TX1	MEAN
DALZ 0104	4.7	4.0	2.7	3.8
GN-Z	4.0	4.3	3.0	3.8
ZORRO	3.7	4.0	2.7	3.4
DALZ 0101	2.7	4.3	3.0	3.3
DALZ 9604	4.0	3.3	2.3	3.2
6186	4.3	4.3	1.0	3.2
HIMENO	.	4.0	2.3	3.2
EMERALD	2.3	4.0	3.0	3.1
DALZ 0105	3.0	3.7	2.3	3.0
DALZ 0102	2.3	3.7	2.7	2.9
BMZ 230	1.3	4.0	2.0	2.4
MEYER	1.0	4.3	1.3	2.2
LSD VALUE	1.3	1.1	1.0	0.7
C.V. (%)	26.9	16.7	25.5	22.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 26A. SEEDHEAD RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2005 DATA

NAME	SEEDHEAD RATINGS 1-9; 9=NONE 2/				
	AR1	MS1	NM1	SC1	MEAN
DALZ 9604	9.0	7.7	8.7	9.0	8.6
6186	9.0	8.0	7.3	9.0	8.3
DALZ 0102	9.0	7.3	7.7	9.0	8.3
DALZ 0104	8.7	8.0	7.3	9.0	8.3
ZORRO	8.7	7.3	7.7	9.0	8.2
DALZ 0101	9.0	7.0	8.3	8.0	8.1
EMERALD	9.0	7.0	8.0	8.3	8.1
GN-Z	8.7	7.3	7.3	9.0	8.1
DALZ 0105	9.0	6.3	8.7	8.0	8.0
HIMENO	4.0	7.7	7.7	9.0	7.1
PZA 32	8.7	2.0	8.0	9.0	6.9
COMPANION	7.7	2.3	8.3	9.0	6.8
PST-R7MA	7.7	3.0	7.7	9.0	6.8
PZB 33	8.3	2.3	8.0	8.3	6.8
CHINESE COMMON	6.7	3.3	7.7	9.0	6.7
ZENITH	6.7	2.3	8.3	9.0	6.6
J-37	7.7	2.3	7.3	9.0	6.6
MEYER	8.0	2.0	7.7	8.3	6.5
PST-R7ZM	7.0	2.3	7.7	9.0	6.5
BMZ 230	5.0	2.7	8.0	9.0	6.2
LSD VALUE	1.8	0.8	1.2	1.1	0.6
C.V. (%)	14.6	10.5	9.1	8.0	11.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 26B. SEEDHEAD RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

NAME	SEEDHEAD RATINGS 1-9; 9=NONE 2/				
	AR1	MS1	NM1	SC1	MEAN
PZA 32	8.7	2.0	8.0	9.0	6.9
COMPANION	7.7	2.3	8.3	9.0	6.8
PST-R7MA	7.7	3.0	7.7	9.0	6.8
PZB 33	8.3	2.3	8.0	8.3	6.8
CHINESE COMMON	6.7	3.3	7.7	9.0	6.7
ZENITH	6.7	2.3	8.3	9.0	6.6
J-37	7.7	2.3	7.3	9.0	6.6
PST-R7ZM	7.0	2.3	7.7	9.0	6.5
LSD VALUE	2.2	0.8	1.3	0.7	0.7
C.V. (%)	17.7	20.0	10.0	4.6	12.5

TABLE 26C. SEEDHEAD RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

NAME	SEEDHEAD RATINGS 1-9; 9=NONE 2/				
	AR1	MS1	NM1	SC1	MEAN
DALZ 9604	9.0	7.7	8.7	9.0	8.6
6186	9.0	8.0	7.3	9.0	8.3
DALZ 0102	9.0	7.3	7.7	9.0	8.3
DALZ 0104	8.7	8.0	7.3	9.0	8.3
ZORRO	8.7	7.3	7.7	9.0	8.2
DALZ 0101	9.0	7.0	8.3	8.0	8.1
EMERALD	9.0	7.0	8.0	8.3	8.1
GN-Z	8.7	7.3	7.3	9.0	8.1
DALZ 0105	9.0	6.3	8.7	8.0	8.0
HIMENO	4.0	7.7	7.7	9.0	7.1
MEYER	8.0	2.0	7.7	8.3	6.5
BMZ 230	5.0	2.7	8.0	9.0	6.2
LSD VALUE	1.6	0.8	1.1	1.4	0.6
C.V. (%)	12.4	8.1	8.5	9.7	10.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 27A. MOWING QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2005 DATA

MOWING QUALITY RATINGS 1-9; 9=CLEAREST CUT 2/

NAME	NM1
EMERALD	6.3
CHINESE COMMON	5.7
J-37	5.7
PST-R7MA	5.3
ZORRO	5.3
6186	5.0
BMZ 230	5.0
COMPANION	5.0
DALZ 0101	5.0
DALZ 0102	5.0
DALZ 0104	5.0
DALZ 0105	5.0
DALZ 9604	5.0
GN-Z	5.0
HIMENO	5.0
MEYER	5.0
PST-R7ZM	5.0
PZA 32	5.0
PZB 33	5.0
ZENITH	5.0
LSD VALUE	0.8
C.V. (%)	9.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 27B. MOWING QUALITY RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

MOWING QUALITY RATINGS 1-9; 9=CLEANEST CUT 2/

NAME	NM1
CHINESE COMMON	5.7
J-37	5.7
PST-R7MA	5.3
COMPANION	5.0
PST-R7ZM	5.0
PZA 32	5.0
PZB 33	5.0
ZENITH	5.0
LSD VALUE	1.0
C.V. (%)	11.8

TABLE 27C. MOWING QUALITY RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

MOWING QUALITY RATINGS 1-9; 9=CLEANEST CUT 2/

NAME	NM1
EMERALD	6.3
ZORRO	5.3
6186	5.0
BMZ 230	5.0
DALZ 0101	5.0
DALZ 0102	5.0
DALZ 0104	5.0
DALZ 0105	5.0
DALZ 9604	5.0
GN-Z	5.0
HIMENO	5.0
MEYER	5.0
LSD VALUE	0.6
C.V. (%)	7.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 28A. MEAN LARGE ZOYSIA PATCH MEASUREMENTS OF ZOYSIAGRASS CULTIVARS 1/  
2005 DATA

MEAN LARGE ZOYSIA PATCH MEASURED IN SQUARE CENTIMETERS 2/

NAME	OK1
ZORRO	3539.0
DALZ 0101	2098.0
DALZ 9604	1927.3
CHINESE COMMON	1844.0
PZB 33	1823.0
HIMENO	1794.3
GN-Z	1712.7
EMERALD	1699.3
ZENITH	1658.0
DALZ 0102	1438.0
J-37	1392.7
COMPANION	1344.0
6186	1256.0
DALZ 0105	983.0
PZA 32	900.7
PST-R7MA	859.0
DALZ 0104	382.5
PST-R7ZM	296.0
MEYER	291.7
BMZ 230	24.3
LSD VALUE	1875.5
C.V. (%)	78.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 28B. MEAN LARGE ZOYSIA PATCH MEASUREMENTS OF ZOYSIAGRASS (SEEDED) CULTIVARS 1/  
2005 DATA

MEAN LARGE ZOYSIA PATCH MEASURED IN SQUARE CENTIMETERS 2/

NAME	OK1
CHINESE COMMON	1844.0
PZB 33	1823.0
ZENITH	1658.0
J-37	1392.7
COMPANION	1344.0
PZA 32	900.7
PST-R7MA	859.0
PST-R7ZM	296.0
LSD VALUE	1510.7
C.V. (%)	72.0

TABLE 28C. MEAN LARGE ZOYSIA PATCH MEASUREMENTS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS 1/  
2005 DATA

MEAN LARGE ZOYSIA PATCH MEASURED IN SQUARE CENTIMETERS 2/

NAME	OK1
ZORRO	3539.0
DALZ 0101	2098.0
DALZ 9604	1927.3
HIMENO	1794.3
GN-Z	1712.7
EMERALD	1699.3
DALZ 0102	1438.0
6186	1256.0
DALZ 0105	983.0
DALZ 0104	382.5
MEYER	291.7
BMZ 230	24.3
LSD VALUE	2129.5
C.V. (%)	82.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 29A. LARGE PATCH RATINGS OF ZOYSIAGRASS CULTIVARS  
AT RALEIGH, NC 1/  
2005 DATA

NAME	LARGE PATCH RATINGS 1-9; 9=NO DISEASE 2/				
	MAY	JUNE	JULY	SEPTEMBER	MEAN
BMZ 230	9.0	9.0	9.0	9.0	9.0
CHINESE COMMON	9.0	9.0	9.0	9.0	9.0
COMPANION	9.0	9.0	9.0	9.0	9.0
DALZ 0101	9.0	9.0	9.0	9.0	9.0
DALZ 0102	9.0	9.0	9.0	9.0	9.0
DALZ 9604	9.0	9.0	9.0	9.0	9.0
EMERALD	9.0	9.0	9.0	9.0	9.0
GN-Z	9.0	9.0	9.0	9.0	9.0
HIMENO	9.0	9.0	9.0	9.0	9.0
PST-R7MA	9.0	9.0	9.0	9.0	9.0
PST-R7ZM	9.0	9.0	9.0	9.0	9.0
PZB 33	9.0	9.0	9.0	9.0	9.0
ZENITH	9.0	9.0	9.0	9.0	9.0
ZORRO	9.0	9.0	9.0	9.0	9.0
6186	8.7	8.3	9.0	9.0	8.8
DALZ 0104	8.7	8.3	9.0	9.0	8.8
J-37	8.7	8.3	9.0	9.0	8.8
PZA 32	8.7	8.3	9.0	9.0	8.8
MEYER	9.0	9.0	9.0	7.7	8.7
DALZ 0105	7.7	6.7	8.7	9.0	8.0
LSD VALUE	0.5	0.9	0.4	0.2	0.3
C.V. (%)	3.3	5.9	1.4	1.4	2.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 29B. LARGE PATCH RATINGS OF ZOYSIAGRASS (SEEDED) CULTIVARS  
AT RALEIGH, NC 1/  
2005 DATA

LARGE PATCH RATINGS 1-9; 9=NO DISEASE 2/					
NAME	MAY	JUNE	JULY	SEPTEMBER	MEAN
CHINESE COMMON	9.0	9.0	9	9	9.0
COMPANION	9.0	9.0	9	9	9.0
PST-R7MA	9.0	9.0	9	9	9.0
PST-R7ZM	9.0	9.0	9	9	9.0
PZB 33	9.0	9.0	9	9	9.0
ZENITH	9.0	9.0	9	9	9.0
J-37	8.7	8.3	9	9	8.8
PZA 32	8.7	8.3	9	9	8.8
LSD VALUE	0.8	1.5	0	0	0.6
C.V. (%)	3.4	6.8	0	0	2.5

TABLE 29C. LARGE PATCH RATINGS OF ZOYSIAGRASS (VEGETATIVE) CULTIVARS  
AT RALEIGH, NC 1/  
2005 DATA

LARGE PATCH RATINGS 1-9; 9=NO DISEASE 2/					
NAME	MAY	JUNE	JULY	SEPTEMBER	MEAN
BMZ 230	9.0	9.0	9.0	9.0	9.0
DALZ 0101	9.0	9.0	9.0	9.0	9.0
DALZ 0102	9.0	9.0	9.0	9.0	9.0
DALZ 9604	9.0	9.0	9.0	9.0	9.0
EMERALD	9.0	9.0	9.0	9.0	9.0
GN-Z	9.0	9.0	9.0	9.0	9.0
HIMENO	9.0	9.0	9.0	9.0	9.0
ZORRO	9.0	9.0	9.0	9.0	9.0
6186	8.7	8.3	9.0	9.0	8.8
DALZ 0104	8.7	8.3	9.0	9.0	8.8
MEYER	9.0	9.0	9.0	7.7	8.7
DALZ 0105	7.7	6.7	8.7	9.0	8.0
LSD VALUE	0.5	0.8	0.4	0.3	0.3
C.V. (%)	3.3	5.3	1.9	1.9	2.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

APPENDIX TABLE. SUMMARY OF TURFGRASS QUALITY RATINGS FOR ZOYSIAGRASS CULTIVARS  
 IN THE 2002 NATIONAL ZOYSIAGRASS TEST \*/  
 2005 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF \*\*/

NAME	QUALITY MEAN 1/	MAXIMUM IN TOP 25% 2/
6186	5.3	6.3
BMZ 230	5.8	31.3
CHINESE COMMON	4.8	0.0
COMPANION	4.8	0.0
DALZ 0101	6.7	75.0
DALZ 0102	6.1	25.0
DALZ 0104	5.9	12.5
DALZ 0105	5.5	12.5
DALZ 9604	5.4	0.0
EMERALD	6.8	87.5
GN-Z	6.3	56.3
HIMENO	6.1	43.8
J-37	4.8	0.0
MEYER	5.6	18.8
PST-R7MA	5.3	0.0
PST-R7ZM	5.2	6.3
PZA 32	5.1	0.0
PZB 33	5.3	6.3
ZENITH	5.3	6.3
ZORRO	6.9	87.5
LSD VALUE	0.2	
C.V. (%)	10.6	

\*/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
 STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

\*\*/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

1/ MEAN - AN AVERAGE OF ALL THE TURFGRASS QUALITY RATINGS FROM ALL LOCATIONS.

2/ MAXIMUM IN TOP 25% - THE PERCENTAGE OF LOCATIONS WHERE THAT ENTRY FINISHED IN THE TOP 25% OF ALL ENTRIES.