

NATIONAL TURFGRASS EVALUATION PROGRAM

The National Turfgrass Evaluation Program (NTEP) is designed to develop and coordinate uniform evaluation trials of turfgrass varieties and promising selections in the United States and Canada. Test results can be used by national companies and plant breeders to determine the broad picture of the adaptation of a cultivar. Results can also be used to determine if a cultivar is well adapted to a local area or level of turf maintenance.

Briefly, the NTEP is a self-supporting, non-profit program, sponsored by the Beltsville Agricultural Research Center and the National Turfgrass Federation, Inc. Program policy is made by a policy committee consisting of one member from each of the four (4) Regional Turfgrass Research Committees in the United States, one member from the Lawn Seed Division of the American Seed Trade Association, one member from the United States Golf Association (USGA) Green Section, one member from the Golf Course Superintendents Assoc. of America (GCSAA), one member for the Turfgrass Producers International (TPI), one member from the Turfgrass Breeders Association and an executive director. The program does not make variety recommendations. However, the data from tests can be used by extension specialists and others for making recommendations.

The policy committee is responsible for determining program policy including, (1) requirements for submission of entries, (2) scheduling tests, (3) evaluation methods, (4) selecting standard or control test entries, (5) setting entry fees, (6) coordinating tests in their respective regions, (7) establishing guidelines for publication and data distribution and (8) scheduling committee meetings.

Executive Director - Kevin N. Morris, National Turfgrass Evaluation Program, Inc.

CURRENT POLICY COMMITTEE MEMBERS:

Dr. Melodee Fraser, Pure-Seed Testing, Inc.
Mr. Paul Hedgpeth, Columbia River Seeds
Dr. Jeff Nus, USGA Green Section
Dr. Michael Richardson, University of Arkansas
Dr. David Kopec, University of Arizona
Mr. Warren Bell, Biograss Sod Farms
Dr. Clark Throssell, Golf Course Superintendents Assoc. of America
Dr. Ken Diesburg, Southern Illinois University
Mr. Pat McClain, Burlingham Seeds
Dr. Scott Ebdon, University of Massachusetts

FOR ADDITIONAL REPORTS OR INFORMATION CONTACT:

Kevin Morris, Executive Director
National Turfgrass Evaluation Program
Beltsville Agricultural Research Center-West
Building 005, Room 307
Beltsville, Maryland 20705
kmorris@ntep.org
www.ntep.org

CONTENTS

2007 National Zoysiagrass Test - 2009 data

LOCATIONS SUBMITTING DATA FOR 2009.....	1
NATIONAL ZOYSIAGRASS TEST, 2007 Entries and Sponsors.....	2
Table A - 2009 Locations, Site Descriptions and Management Practices in the 2007 National Zoysiagrass Test.....	3
Table B - Locations and Data Collected in 2009.....	3
Table 1 - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars at Riverside, CA.....	4
Table 2 - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars at Gainesville, FL.....	4
Table 3 - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars at West Lafayette, IN.....	5
Table 4 - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars at Manhattan, KS.....	5
Table 5 - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars at Raleigh, NC.....	6
Table 6 - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars at Dallas, TX.....	6
Table 7 - Percent Establishment and Other Ratings of Zoysiagrass Cultivars Grown under Salt Stress at Las Cruces, NM.....	7
Table 8 - Genetic Color Ratings of Zoysiagrass Cultivars.....	7
Table 9 - Spring Greenup Ratings of Zoysiagrass Cultivars.....	8
Table 10- Leaf Texture Ratings of Zoysiagrass Cultivars.....	8
Table 11- Spring Density Ratings of Zoysiagrass Cultivars.....	9
Table 12- Summer Density Ratings of Zoysiagrass Cultivars.....	9
Table 13- Fall Density Ratings of Zoysiagrass Cultivars.....	10
Table 14- Percent Living Ground Cover (Spring) Ratings of Zoysiagrass Cultivars.....	10
Table 15- Percent Living Ground Cover (Summer) Ratings of Zoysiagrass Cultivars.....	11
Table 16- Winter Color Ratings of Zoysiagrass Cultivars.....	11

CONTENTS (continued)

Table 17- Percent Winter Kill Ratings of Zoysiagrass Cultivars.....	12
Table 18- Brown Patch (Cool Temperature) Ratings of Zoysiagrass Cultivars.....	12
Table 19- Fall Color (September) Ratings of Zoysiagrass Cultivars.....	13
Table 20- Fall Color (October) Ratings of Zoysiagrass Cultivars.....	13
Table 21- Fall Color (November) Ratings of Zoysiagrass Cultivars.....	14
Table 22- Fall Color (December) Ratings of Zoysiagrass Cultivars.....	14
Table 23- Seedhead Ratings of Zoysiagrass Cultivars.....	15
Table 24- Mole Cricket Ratings of Zoysiagrass Cultivars.....	15
Table 25- Dollar Spot Ratings of Zoysiagrass Cultivars at Raleigh, NC.....	16

A Guide to NTEP Turfgrass Ratings

Introduction

The quality and scientific merit of NTEP data is extremely important. However, the evaluation of turfgrass species and cultivars is a difficult and complex issue. Furthermore, turfgrass evaluation is generally a subjective process based on visual estimates of factors, like genetic color, stand density, leaf texture, uniformity and quality. These factors can not be measured in the same way as other agricultural crops. Turfgrass quality is not a measure of yield or nutritive value. Turfgrass quality is a measure of aesthetics (i.e. density, uniformity, texture, smoothness, growth habit and color), and functional use. The most common way of assessing turfgrass quality is a visual rating system that is based on the turfgrass evaluator's judgement.

General Considerations

Most visual ratings collected on NTEP trials are based on a 1 to 9 rating scale. One is the poorest or lowest and 9 is the best or highest rating. However, a few characteristics, such as winter kill or percent living ground cover, are rated on a percentage basis, again by using the evaluator's judgement. Most disease ratings found in NTEP reports will use the 1-9 scale, 9=no disease except where the evaluator made a judgement of the percentage of disease in each plot. Percent disease data will be found in separate tables and will normally not be included with disease data using the 1-9 scale.

Turfgrass Quality

Turfgrass Quality is based on 9 being outstanding or ideal turf and 1 being poorest or dead. A rating of 6 or above is generally considered acceptable. A quality rating value of 9 is reserved for a perfect or ideal grass, but it also can reflect an absolutely outstanding treatment plot. The NTEP requires quality ratings on a monthly basis. Quality ratings take into account the aesthetic and functional aspects of the turf. Quality ratings are not based on color alone, but on a combination of color, density, uniformity, texture, and disease or environmental stress.

Turfgrass quality ratings are grouped and presented by region, management level, a particular stress (shade, traffic, etc.) and in some cases, by individual location (starting with 2007 data, data from each location will be posted separately as well on the NTEP web site, <http://www.ntep.org>). Also available now is a summary table (Appendix) in the back of this report. This summary table includes various statistical measures not previously compiled for NTEP reports. For an explanation of this table and these changes, please go to the NTEP web site at <http://www.ntep.org/pdf/grandmean.mem.pdf>.

Other Ratings

More detailed information on the ratings of specific characteristics can be found on the NTEP web site at <http://www.ntep.org/reports/ratings.htm>.

2007 NATIONAL ZOYSIAGRASS TEST

LOCATIONS SUBMITTING DATA FOR 2009

<u>State</u>	<u>Location</u>	<u>Code</u>
California	Riverside	CA3
Florida	Gainesville	FL1
Indiana	West Lafayette	IN1
Kansas	Manhattan	KS1
New Mexico	Las Cruces (Salt Tolerance)	NM1
North Carolina	Raleigh	NC1
Texas	Dallas	TX1

2007 NATIONAL ZOYSIAGRASS TEST

Entries and Sponsors

Entry No.	Name	Type	Sponsor
* 1	Zenith	seeded	Standard entry
* 2	Meyer	vegetative	Standard entry
* 3	Zorro	vegetative	Standard entry
4	DALZ 0501	vegetative	Texas A&M Dallas & Phillip Jennings Turf
5	DALZ 0701	vegetative	Texas A&M Dallas
6	DALZ 0702	vegetative	Texas A&M Dallas
* 7	Shadowturf	vegetative	Ivey Gardens
8	L1F	vegetative	Bladerunner Farms
9	29-2	vegetative	Bladerunner Farms
10	240	vegetative	Bladerunner Farms
11	380-1	vegetative	Bladerunner Farms

* COMMERCIALIALLY AVAILABLE IN THE USA IN 2010.

TABLE A.

2009 LOCATIONS, SITE DESCRIPTIONS AND MANAGEMENT PRACTICES IN
THE 2002 NATIONAL ZOYSIAGRASS TEST

LOCATION	SOIL TEXTURE	SOIL PH	SOIL	SOIL	NITROGEN (LBS/1000 SQ FT)	SUN OR SHADE	MOWING HEIGHT (IN)	IRRIGATION PRACTICED
			PHOSPHOROUS (LBS/ACRE)	POTASSIUM (LBS/ACRE)				
CA3	SANDY LOAM	7.1-7.5	-	-	-	FULL SUN	1.6-2.0	TO PREVENT STRESS
FL1	SAND	6.6-7.0	-	-	0.0-1.0	FULL SUN	1.6-2.0	TO PREVENT STRESS
IN1	SANDY CLAY LOAM	7.1-7.5	61-150	241-375	2.1-3.0	FULL SUN	0.6-1.0	-
KS1	-	-	-	-	-	-	-	-
NC1	SILTY CLAY AND CLAY	6.1-6.5	61-150	0-150	3.1-4.0	FULL SUN	2.1-2.5	TO PREVENT STRESS
NM1	LOAMY SAND	7.6-8.5	-	-	5.1-6.0	FULL SUN	0.6-1.0	TO PREVENT STRESS
TX1	SILTY CLAY AND CLAY	7.6-8.5	151-270	241-375	3.1-4.0	FULL SUN	1.6-2.0	TO PREVENT DORMANCY

TABLE B.

LOCATIONS AND DATA COLLECTED IN 2009

	JANUARY	FEBRUARY	MARCH	APRIL	MAY	JUNE	JULY	AUGUST	SEPTEMBER	OCTOBER	NOVEMBER	DECEMBER			
LOCATION	QUALITY RATING	GENETIC COLOR	SPRING GREENUP	LEAF TEXTURE											
CA3			X	X	X	X	X	X	X	X	X	X	X	X	X
FL1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
IN1					X	X	X	X	X	X	X	X	X	X	X
KS1					X	X	X	X					X	X	X
NC1				X	X	X	X	X	X	X	X	X	X	X	X
NM1					X	X	X	X	X	X	X	X	X	X	X
TX1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X

TABLE B. (CONT'D)

LOCATIONS AND DATA COLLECTED IN 2009

LOCATION	SPRING DENSITY	SUMMER DENSITY	FALL DENSITY	PERCENT COVER SPRING	PERCENT COVER SUMMER	PERCENT COVER FALL	PERCENT WINTER COLOR	PERCENT WINTER KILL	BROWN PATCH COOL TEMP.	FALL COLOR SEPTEMBER	FALL COLOR OCTOBER	FALL COLOR NOVEMBER	FALL COLOR DECEMBER	FALL SEEDHEAD RATINGS	MOLE CRICKET BILLBUG	PERCENT DOLLAR JULY	SPOT OCT
CA3						X								X			
FL1	X	X	X	X	X	X	X	X	X	X	X	X	X	X			
IN1								X									
KS1					X						X				X		
NC1												X		X			
NM1					X	X	X	X			X	X	X	X		X	
TX1	X	X	X									X	X				

TABLE 1. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS
AT RIVERSIDE, CA 1/
2009 DATA

NAME	TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/									
	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	MEAN
* ZORRO	7.0	6.7	7.3	6.7	7.3	8.3	7.7	8.0	7.0	7.3
DALZ 0701	6.3	6.0	6.0	6.0	6.7	7.0	6.7	8.3	6.7	6.6
DALZ 0501	6.7	6.3	6.3	6.0	6.3	5.7	5.3	7.7	5.7	6.2
DALZ 0702	6.3	6.7	6.7	5.7	6.0	6.0	6.0	8.0	4.7	6.2
L1F	6.3	6.3	6.7	5.7	6.0	6.3	5.0	8.0	5.3	6.2
* SHADOWTURF	6.3	6.7	6.3	5.7	6.7	5.7	3.7	7.7	3.0	5.7
29-2	5.3	7.0	5.0	5.0	5.7	6.0	4.7	5.3	4.7	5.4
* ZENITH	5.0	6.7	4.3	4.0	5.3	6.0	4.3	5.0	4.0	5.0
380-1	4.7	5.7	5.0	3.0	3.3	4.7	3.7	6.0	3.7	4.4
240	3.0	5.7	3.3	3.0	3.7	5.0	4.0	4.0	2.7	3.8
* MEYER	4.3	5.3	4.3	2.3	2.7	4.3	3.3	4.7	2.3	3.7
LSD VALUE	1.4	1.0	1.2	1.3	1.2	1.9	1.3	0.8	1.3	0.9
C.V. (%)	14.9	8.5	13.4	16.5	14.0	17.1	15.6	7.6	18.2	10.0

TABLE 2. MEAN TURFGRASS QUALITY AND RATINGS OF ZOYSIAGRASS CULTIVARS
GAINESVILLE, FL 1/
2009 DATA

NAME	TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/												
	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	MEAN
DALZ 0701	6.0	4.3	4.0	5.0	4.0	5.3	5.0	5.0	5.7	7.3	8.3	5.3	5.4
L1F	6.3	4.3	4.3	4.7	4.0	4.7	5.3	5.7	5.3	7.0	7.3	5.7	5.4
SHADOWTURF	6.3	4.0	4.3	5.0	4.0	5.0	5.0	5.3	5.7	6.3	6.7	5.7	5.3
DALZ 0702	5.3	3.7	3.3	4.3	4.0	5.3	4.7	6.0	3.7	6.7	7.0	6.3	5.0
DALZ 0501	5.7	3.7	3.0	4.3	2.7	4.3	4.3	3.3	5.3	6.3	7.0	7.3	4.8
ZORRO	4.3	3.0	3.0	4.0	2.3	3.3	4.0	4.7	5.0	5.0	5.7	5.0	4.1
ZENITH	3.3	2.0	2.3	2.7	3.0	5.0	5.7	5.3	4.7	4.0	3.0	2.3	3.6
240	2.7	1.0	1.3	2.3	2.7	3.7	4.7	4.0	4.3	3.0	3.3	3.3	3.0
29-2	4.0	2.0	2.0	2.0	2.7	3.7	2.7	3.3	2.7	3.3	3.0	2.3	2.8
380-1	2.7	1.3	2.3	3.0	2.3	4.0	2.3	4.0	3.0	2.0	1.0	1.3	2.4
MEYER	3.0	1.0	1.0	1.3	2.0	3.3	3.0	2.3	2.3	2.3	2.3	1.7	2.1
LSD VALUE	1.1	0.7	1.2	1.9	1.7	2.6	1.8	3.2	1.7	1.6	1.6	1.4	0.9
C.V. (%)	15.2	16.9	26.2	30.2	27.0	25.5	22.7	32.2	22.2	20.0	20.3	20.4	14.6

* COMMERCIALLY AVAILABLE IN THE USA IN 2010.

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 3. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS
AT WEST LAFAYETTE, IN 1/
2009 DATA

	TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/						
NAME	MAY	JUN	JUL	AUG	SEP	OCT	MEAN
29-2	6.0	7.7	6.0	7.0	7.7	6.3	6.8
MEYER	6.7	7.3	7.3	7.0	6.7	5.7	6.8
ZENITH	6.7	7.7	5.7	6.7	7.3	6.7	6.8
240	5.3	6.3	6.0	5.7	6.7	5.7	5.9
ZORRO	2.0	2.0	3.0	3.0	4.0	2.7	2.8
380-1	2.3	2.0	2.7	3.0	2.7	2.7	2.6
DALZ 0701	1.0	2.0	2.0	2.0	3.0	2.0	2.0
DALZ 0501
DALZ 0702
L1F
SHADOWTURF
LSD VALUE	2.0	1.9	1.7	1.9	1.5	1.8	1.1
C.V. (%)	19.9	18.2	17.2	17.8	13.6	18.5	11.6

TABLE 4. MEAN TURFGRASS QUALITY AND OTHER RATINGS OF ZOYSIAGRASS CULTIVARS
IN THE 2007 NATIONAL ZOYSIAGRASS TEST AT MANHATTAN, KS 1/
2009 DATA

	TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/				
NAME	MAY	JUN	JUL	AUG	MEAN
380-1	4.3	7.0	8.7	8.7	7.2
DALZ 0701	2.7	8.0	8.0	8.3	6.8
ZORRO	4.0	6.7	8.0	8.3	6.8
MEYER	5.3	6.7	7.0	7.0	6.5
29-2	5.7	6.3	6.7	7.0	6.4
240	4.7	7.0	7.0	6.7	6.3
DALZ 0702	1.7	6.7	7.7	8.0	6.0
ZENITH	6.3	5.3	6.0	6.3	6.0
DALZ 0501	2.3	5.7	7.3	8.0	5.8
SHADOWTURF	1.0	4.0	5.3	7.0	4.3
L1F	1.0	1.0	1.0	1.0	1.0
LSD VALUE	1.5	1.2	1.2	1.1	0.7
C.V. (%)	26.4	12.6	11.2	10.5	8.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 5. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS
AT RALEIGH, NC 1/
2009 DATA

NAME	TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/									
	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	MEAN	
380-1	7.3	8.0	8.3	8.3	8.7	8.0	7.3	7.7	8.0	
ZORRO	6.7	7.3	6.7	6.7	7.7	7.3	6.7	7.7	7.1	
MEYER	6.3	6.3	7.0	7.7	7.7	7.0	6.7	7.0	7.0	
DALZ 0702	6.3	7.0	6.0	6.0	6.7	6.7	7.0	7.7	6.7	
DALZ 0501	4.3	4.7	7.0	7.0	7.0	8.0	7.7	7.3	6.6	
29-2	5.3	5.7	6.7	6.3	7.0	7.0	5.7	6.0	6.2	
240	4.7	5.3	7.0	6.3	7.0	7.0	5.7	5.3	6.0	
DALZ 0701	5.3	6.0	4.7	5.0	6.7	6.7	6.0	7.0	5.9	
ZENITH	5.0	5.7	6.7	6.0	5.7	5.7	6.0	5.7	5.8	
SHADOWTURF	4.7	4.7	5.7	6.0	6.0	6.7	6.0	5.7	5.7	
L1F	4.0	4.0	5.3	6.0	6.7	6.3	5.7	5.7	5.5	
LSD VALUE	1.3	1.2	1.3	1.3	1.3	1.0	1.7	1.3	0.8	
C.V. (%)	14.1	12.4	11.4	11.0	10.2	7.9	12.9	11.2	7.8	

TABLE 6. MEAN TURFGRASS QUALITY AND OTHER RATINGS OF ZOYSIAGRASS CULTIVARS
IN THE 2007 NATIONAL ZOYSIAGRASS TEST AT DALLAS, TX 1/
2009 DATA

NAME	TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/												
	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	MEAN
ZORRO	6.0	5.0	6.3	5.7	7.0	7.0	7.7	8.0	7.0	8.0	7.3	7.3	6.9
DALZ 0702	6.0	5.0	5.7	6.0	7.0	6.7	8.0	7.3	7.0	7.7	6.7	7.0	6.7
DALZ 0701	6.0	5.0	6.3	6.0	7.0	7.0	7.0	6.7	7.0	7.3	7.3	7.0	6.6
DALZ 0501	6.0	5.0	6.7	5.0	6.7	6.0	7.0	7.0	6.7	7.0	7.7	7.0	6.5
L1F	5.7	4.7	6.3	5.0	6.3	6.0	7.0	7.0	7.0	7.3	8.0	7.0	6.4
380-1	4.3	3.3	6.0	7.0	7.0	7.3	7.3	7.3	6.3	7.7	6.7	5.3	6.3
SHADOWTURF	5.0	4.3	5.3	5.3	6.3	6.3	7.0	7.3	6.7	7.3	7.7	7.0	6.3
240	4.3	3.3	5.3	5.7	5.7	5.3	6.0	6.0	5.0	5.7	5.7	5.0	5.3
29-2	3.7	2.3	5.0	4.7	4.7	5.0	6.0	6.0	5.3	6.0	5.7	4.3	4.9
MEYER	4.0	2.3	5.3	5.0	4.7	4.3	5.7	5.3	4.7	5.3	5.0	4.3	4.7
ZENITH	4.0	2.3	5.0	5.0	4.3	4.0	5.7	5.7	5.0	5.7	5.7	4.0	4.7
LSD VALUE	0.9	1.1	1.4	3.1	1.9	2.1	1.8	1.3	1.2	1.0	0.9	1.0	0.9
C.V. (%)	10.9	17.1	11.8	21.5	16.3	18.2	12.8	10.6	11.3	8.8	8.4	10.6	9.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 7.

MEAN TURFGRASS QUALITY AND OTHER RATINGS OF ZOYSIAGRASS CULTIVARS 1/
GROWN UNDER SALT STRESS AT LAS CRUCES, NM */
2009 DATA

NAME	GENETIC COLOR	SPRING GREENUP	TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/									QUALITY RATINGS				
			PERCENT COVER SPRING	PERCENT COVER SUMMER	PERCENT COVER FALL	FALL COLOR SEPTEMBER	FALL COLOR OCTOBER	FALL COLOR NOVEMBER	JUL	AUG	SEP	OCT	MEAN			
SHADOWTURF	8.3	3.3	80.3	99.0	99.0	7.7	5.7	1.7	7.7	8.3	8.0	7.7	7.9			
L1F	8.3	3.0	78.7	98.7	99.0	8.0	7.0	2.3	7.0	8.0	8.0	7.3	7.6			
DALZ 0701	7.7	3.3	79.0	84.7	92.7	7.0	6.0	3.0	6.3	7.0	6.7	6.3	6.6			
DALZ 0501	6.3	2.0	51.3	63.3	91.3	7.0	6.7	2.7	5.3	6.3	6.7	6.0	6.1			
380-1	7.7	3.7	66.0	86.3	80.0	7.3	5.7	2.0	5.3	6.7	5.7	5.7	5.8			
DALZ 0702	6.7	2.7	60.0	88.0	96.0	7.0	6.0	2.0	6.0	6.0	5.3	5.7	5.8			
ZORRO	5.0	1.3	19.3	25.7	46.3	4.7	4.3	2.0	4.3	5.0	4.7	5.0	4.8			
240	5.7	2.0	29.7	30.0	21.7	3.7	3.3	1.3	4.3	4.3	4.0	4.0	3.7	4.1		
29-2	4.7	2.3	28.0	23.0	22.0	3.7	3.3	2.0	3.7	4.0	3.7	3.3	3.7			
ZENITH	4.3	1.3	18.0	19.7	21.3	3.3	2.7	1.7	3.3	3.3	3.3	3.0	3.3			
MEYER	4.7	1.7	7.0	14.3	15.0	2.7	2.3	1.3	2.0	2.3	2.3	3.0	2.4			
LSD VALUE	0.7	0.8	27.2	19.9	22.2	1.3	1.3	0.8	1.5	1.3	1.2	1.3	1.1			
C.V. (%)	7.1	19.3	35.1	22.1	22.7	14.6	17.2	21.8	18.6	15.2	14.5	15.6	12.8			

*/ THE FOLLOWING IS INFORMATION ON SALINITY AT THIS SITE:

WATER QUALITY: PH 7.9, TOTAL DISSOLVED SOLIDS 1800, SAR 2.06

SOIL QUALITY (MEASURED IN JUNE, NOV.; 0-4 INCH DEPTH): PH 7.8, 7.8; EC (dS/m) 2.1, 5.2; SAR 2.1, 8.5

SOIL QUALITY (MEASURED IN JUNE, NOV.; 4-8 INCH DEPTH): PH 8.4, 8.0; EC (dS/m) 1.6, 2.4; SAR 2.2, 4.1

TABLE 8. GENETIC COLOR RATINGS OF ZOYSIAGRASS CULTIVARS 1/
2009 DATA
GENETIC COLOR RATINGS 1-9; 9=DARK GREEN 2/

NAME	CA3	FL1	IN1	KS1	NC1	TX1	MEAN
380-1	5.3	7.7	8.7	8.0	8.0	9.0	7.8
29-2	7.3	7.0	8.0	6.7	6.7	8.0	7.3
240	7.3	7.0	8.0	7.7	7.0	6.3	7.2
MEYER	5.3	6.7	7.3	7.0	6.3	7.3	6.7
ZORRO	6.7	6.3	7.7	7.0	4.0	7.7	6.6
ZENITH	7.3	6.3	7.0	7.0	5.7	5.7	6.5
DALZ 0501	5.7	6.7	.	7.3	6.0	6.7	6.5
SHADOWTURF	5.0	7.7	.	7.0	5.0	7.3	6.4
DALZ 0702	5.7	6.7	.	7.0	5.0	7.3	6.3
DALZ 0701	4.7	7.3	6.0	7.3	4.0	8.0	6.2
L1F	5.0	7.0	.	.	4.3	7.7	6.0
LSD VALUE	1.7	1.0	1.1	0.6	1.0	1.7	0.5
C.V. (%)	17.6	9.4	7.7	5.1	11.1	14.0	11.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 9. SPRING GREENUP RATINGS OF ZOYSIAGRASS CULTIVARS 1/
2009 DATA

SPRING GREENUP RATINGS 1-9; 9=COMPLETELY GREEN 2/

NAME	CA3	FL1	IN1	KS1	NC1	TX1	MEAN
DALZ 0701	5.3	6.3	1.0	.	5.0	5.7	4.7
DALZ 0702	5.0	5.7	1.0	.	5.0	5.3	4.4
L1F	6.0	6.0	1.0	.	3.3	5.7	4.4
29-2	1.3	4.7	4.0	5.3	5.3	5.0	4.3
SHADOTURF	5.7	6.3	1.0	.	2.7	5.0	4.1
ZENITH	1.3	3.0	4.0	5.0	5.7	5.7	4.1
380-1	2.7	3.3	1.3	2.0	6.7	8.0	4.0
DALZ 0501	5.7	4.7	1.0	.	2.3	6.3	4.0
MEYER	2.3	2.0	3.3	2.0	6.0	5.7	3.6
ZORRO	4.0	3.7	1.0	1.0	4.3	7.0	3.5
240	1.0	1.7	2.7	2.3	6.0	6.3	3.3
LSD VALUE	0.7	1.3	0.5	0.8	1.1	1.7	0.5
C.V. (%)	12.6	18.1	15.5	17.9	14.6	17.7	17.3

TABLE 10. LEAF TEXTURE RATINGS OF ZOYSIAGRASS CULTIVARS 1/
2009 DATA

LEAF TEXTURE RATINGS 1-9; 9=VERY FINE 2/

NAME	CA3	FL1	IN1	KS1	NC1	TX1	MEAN
DALZ 0501	8.3	8.0	.	9.0	8.3	7.3	8.2
SHADOTURF	9.0	8.0	.	9.0	8.7	6.3	8.2
L1F	8.7	8.0	.	.	8.7	7.3	8.2
ZORRO	8.3	8.0	9.0	9.0	8.0	6.7	8.2
DALZ 0702	8.7	8.0	.	9.0	8.0	7.0	8.1
380-1	7.0	8.0	8.3	9.0	7.7	6.3	7.7
DALZ 0701	8.0	8.3	5.0	9.0	8.0	7.0	7.6
MEYER	5.3	6.0	7.0	6.7	7.0	5.7	6.3
29-2	5.3	6.0	7.0	7.0	5.7	5.3	6.1
240	4.3	6.0	6.0	7.0	5.7	5.7	5.8
ZENITH	4.0	5.7	5.3	5.3	5.0	5.0	5.1
LSD VALUE	0.9	0.6	0.6	0.4	0.7	1.0	0.3
C.V. (%)	7.9	5.4	4.9	3.2	5.8	10.3	6.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 11. SPRING DENSITY RATINGS OF ZOYSIAGRASS CULTIVARS 1/
2009 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	FL1	TX1	MEAN
DALZ 0701	6.0	6.7	6.3
SHADOWTURF	6.7	5.7	6.2
DALZ 0702	4.7	7.0	5.8
L1F	5.7	5.7	5.7
DALZ 0501	4.7	6.3	5.5
ZORRO	4.0	6.7	5.3
380-1	2.3	8.0	5.2
ZENITH	2.3	6.0	4.2
240	1.7	6.0	3.8
29-2	1.7	6.0	3.8
MEYER	2.0	5.7	3.8
LSD VALUE	2.2	2.1	1.5
C.V. (%)	36.2	20.2	26.2

TABLE 12. SUMMER DENSITY RATINGS OF ZOYSIAGRASS CULTIVARS 1/
2009 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	FL1	NC1	TX1	MEAN
DALZ 0702	5.0	6.3	9.0	6.8
DALZ 0501	4.0	6.7	9.0	6.6
L1F	5.0	5.7	9.0	6.6
ZORRO	4.0	6.7	9.0	6.6
SHADOWTURF	5.0	5.7	8.7	6.4
DALZ 0701	4.7	5.7	8.7	6.3
ZENITH	6.0	6.3	6.7	6.3
380-1	2.3	7.3	8.3	6.0
240	4.3	5.7	6.7	5.6
29-2	3.0	5.7	7.0	5.2
MEYER	3.0	6.7	6.0	5.2
LSD VALUE	2.0	1.8	1.5	1.0
C.V. (%)	29.5	17.9	11.7	18.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 13. FALL DENSITY RATINGS OF ZOYSIAGRASS CULTIVARS 1/
2009 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME	FL1	TX1	MEAN
DALZ 0701	8.3	9.0	8.7
DALZ 0501	7.3	9.0	8.2
L1F	7.3	9.0	8.2
DALZ 0702	7.0	9.0	8.0
SHADOWTURF	7.0	9.0	8.0
ZORRO	7.0	9.0	8.0
29-2	4.3	9.0	6.7
ZENITH	4.3	8.3	6.3
240	3.3	7.7	5.5
MEYER	3.7	7.3	5.5
380-1	1.3	9.0	5.2
LSD VALUE	1.8	1.1	1.0
C.V. (%)	20.1	7.8	13.0

TATLE 14. PERCENT LIVING GROUND COVER (SPRING) RATINGS OF ZOYSIAGRASS CULTIVARS 1/
2009 DATA

PERCENT LIVING GROUND COVER IN SPRING: LOCATIONS 2/

NAME	FL1	KS1	MEAN
29-2	50.0	88.3	69.2
ZENITH	38.3	90.0	64.2
MEYER	25.0	86.7	55.8
380-1	38.3	65.0	51.7
240	15.0	88.3	51.7
DALZ 0701	73.3	26.7	50.0
ZORRO	48.3	38.3	43.3
DALZ 0702	66.7	18.3	42.5
DALZ 0501	65.0	18.3	41.7
SHADOWTURF	75.0	3.0	39.0
L1F	73.3	0.0	36.7
LSD VALUE	13.7	19.6	12.0
C.V. (%)	16.5	25.7	21.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 15. PERCENT LIVING GROUND COVER (SUMMER) RATINGS OF ZOYSIAGRASS CULTIVARS 1/
2009 DATA

PERCENT LIVING GROUND COVER IN SUMMER: LOCATIONS 2/

NAME	FL1
ZENITH	98.0
L1F	93.3
SHADOWTURF	93.3
240	90.3
DALZ 0701	90.0
DALZ 0702	90.0
MEYER	90.0
29-2	88.3
DALZ 0501	88.3
ZORRO	86.7
380-1	76.7
LSD VALUE	10.0
C.V. (%)	7.0

TABLE 16. WINTER COLOR RATINGS OF ZOYSIAGRASS CULTIVARS 1/
2009 DATA

WINTER COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	CA3	FL1	NC1	MEAN
DALZ 0501	6.3	7.0	4.7	6.0
DALZ 0702	5.0	6.7	4.7	5.4
L1F	4.7	7.0	4.3	5.3
SHADOWTURF	4.7	7.0	4.3	5.3
DALZ 0701	6.0	5.3	4.0	5.1
ZORRO	3.7	4.3	6.0	4.7
29-2	1.7	3.3	7.7	4.2
ZENITH	1.3	2.7	8.3	4.1
MEYER	1.0	2.7	8.7	4.1
240	1.0	3.0	7.7	3.9
380-1	1.7	1.0	7.0	3.2
LSD VALUE	1.8	1.6	1.3	0.9
C.V. (%)	32.7	22.0	13.6	21.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 17. PERCENT WINTER KILL RATINGS OF ZOysiAGRASS CULTIVARS 1/
2009 DATA

PERCENT WINTER KILL RATINGS: LOCATIONS 2/

NAME	IN1
DALZ 0501	99.0
DALZ 0702	99.0
L1F	99.0
SHADOWTURF	99.0
DALZ 0701	97.3
380-1	83.0
ZORRO	80.7
240	22.3
29-2	4.0
ZENITH	2.7
MEYER	1.7
LSD VALUE	13.0
C.V. (%)	12.9

TABLE 18. BROWN PATCH (WARM TEMPERATURE) RATINGS OF ZOysiAGRASS CULTIVARS 1/
2009 DATA

BROWN PATCH RATINGS 1-9; 9=NO DISEASE 2/

NAME	FL1
DALZ 0501	7.7
L1F	7.7
DALZ 0702	7.3
SHADOWTURF	7.3
ZORRO	6.7
DALZ 0701	6.3
240	3.7
29-2	3.3
380-1	3.3
MEYER	3.0
ZENITH	3.0
LSD VALUE	1.8
C.V. (%)	20.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 19. FALL COLOR (SEPTEMBER) RATINGS OF ZOYSIAGRASS CULTIVARS 1/
2009 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	FL1
DALZ 0701	6.3
29-2	6.0
SHADOWTURF	5.7
240	5.3
DALZ 0501	5.0
L1F	5.0
DALZ 0702	4.7
380-1	4.3
MEYER	4.3
ZENITH	4.3
ZORRO	4.3
LSD VALUE	2.7
C.V. (%)	33.7

TABLE 20. FALL COLOR (OCTOBER) RATINGS OF ZOYSIAGRASS CULTIVARS 1/
2009 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	FL1	KS1	MEAN
DALZ 0701	7.7	7.0	7.3
L1F	7.3	.	7.3
DALZ 0702	7.3	7.0	7.2
29-2	7.0	6.7	6.8
SHADOWTURF	7.0	6.7	6.8
DALZ 0501	6.3	6.7	6.5
ZORRO	4.7	6.7	5.7
ZENITH	4.3	5.0	4.7
240	4.0	5.0	4.5
380-1	3.7	5.3	4.5
MEYER	2.3	6.3	4.3
LSD VALUE	2.6	1.0	1.5
C.V. (%)	28.8	10.1	21.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 21. FALL COLOR (NOVEMBER) RATINGS OF ZOYSIAGRASS CULTIVARS 1/
2009 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	FL1	NC1	TX1	MEAN
DALZ 0702	7.7	7.7	7.0	7.4
DALZ 0701	7.0	7.7	7.0	7.2
380-1	8.0	6.3	6.3	6.9
L1F	7.3	6.0	7.3	6.9
DALZ 0501	5.0	7.3	7.3	6.6
SHADOWTURF	7.0	5.7	6.3	6.3
ZORRO	5.3	5.3	6.7	5.8
29-2	7.7	4.0	5.3	5.7
240	6.0	3.3	5.7	5.0
MEYER	5.0	3.3	5.7	4.7
ZENITH	4.7	2.3	5.0	4.0
LSD VALUE	1.5	1.0	1.0	0.7
C.V. (%)	14.6	11.2	10.3	12.3

TABLE 22. FALL COLOR (DECEMBER) RATINGS OF ZOYSIAGRASS CULTIVARS 1/
2009 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME	CA3	FL1	TX1	MEAN
DALZ 0501	5.0	8.3	7.3	6.9
L1F	4.7	7.3	7.3	6.4
DALZ 0702	4.7	7.0	7.0	6.2
DALZ 0701	5.0	5.7	7.0	5.9
SHADOWTURF	4.0	6.7	6.3	5.7
ZORRO	3.7	5.3	6.7	5.2
240	2.0	3.3	5.7	3.7
29-2	2.0	3.3	5.3	3.6
380-1	2.3	2.0	6.3	3.6
MEYER	2.0	2.3	5.7	3.3
ZENITH	2.0	3.0	5.0	3.3
LSD VALUE	0.7	1.5	1.0	0.7
C.V. (%)	13.6	19.0	10.3	14.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 23. SEEDHEAD RATINGS OF ZOYSIAGRASS CULTIVARS 1/
2009 DATA

SEEDHEAD RATINGS 1-9; 9=NONE 2/

NAME	NC1
380-1	9.0
DALZ 0501	9.0
DALZ 0701	9.0
DALZ 0702	9.0
L1F	9.0
SHADOWTURF	9.0
ZORRO	9.0
ZENITH	6.7
MEYER	6.0
29-2	4.3
240	3.7
LSD VALUE	0.5
C.V. (%)	4.0

TABLE 24. MOLE CRICKET RATINGS OF ZOYSIAGRASS CULTIVARS 1/
2009 DATA

MOLE CRICKET RATINGS 1-9; 9=NO DAMAGE 2/

NAME	FL1
380-1	8.0
MEYER	8.0
240	7.7
ZENITH	7.7
L1F	7.3
29-2	7.0
SHADOWTURF	7.0
DALZ 0501	6.7
ZORRO	6.3
DALZ 0702	6.0
DALZ 0701	5.7
LSD VALUE	1.4
C.V. (%)	12.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 25. DOLLAR SPOT RATINGS OF ZOYSIAGRASS CULTIVARS
AT RALEIGH, NC 1/
2009 DATA

DOLLAR SPOT RATINGS 1-9; 9=NO DISEASE 2/

NAME	JULY	OCTOBER	MEAN
MEYER	9.0	8.0	8.5
240	9.0	7.3	8.2
29-2	9.0	7.3	8.2
ZENITH	9.0	7.3	8.2
380-1	8.7	7.0	7.8
DALZ 0501	6.3	7.3	6.8
SHADOWTURF	6.0	7.3	6.7
ZORRO	5.7	7.0	6.3
L1F	5.0	7.0	6.0
DALZ 0702	4.0	5.0	4.5
DALZ 0701	3.7	4.7	4.2
LSD VALUS	1.3	1.6	0.8
C.V. (%)	12.2	13.1	7.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.