

TABLE 2A. 2000 MANAGEMENT - ON-SITE BERMUDAGRASS TEST AT MOBILE, AL (COUNTRY CLUB OF MOBILE)

ESTABLISHMENT		FERTILIZATION			HERBICIDES		
Planting date	18-Jun-98	Date(s)	Product	Rate (lbs./M)	Date(s)	Product	Rate (oz./M)
Problems during	None						
	1 lb. N/M, amend soil with Ca as needed. Topdress & roll until smooth surface was attained	June - October November - May		1 lb. N & 2 lbs. K 1/2 lb. N & 1 lb. K	None		
				75% slow release (Nutralene) Nitrogen Typical blend 12-2-24 Foliar nutrients as needed (<2 lbs./year)			
FACTORS OF PLAY							
Date opened for play		FUNGICIDES			INSECTICIDES		
Date closed for play	NA	Date(s)	Product	Rate (oz./M)	Date(s)	Product	Rate (oz./M)
Type of spikes allowed	softspikes						
Uses of green	putting, chipping,	July-September Fall, Spring	Delta Guard for army worms Banol, Heritage, Terrazole for Pythium		None		
MOWING							
Initial height	0.155"	Chipco, chlorothalonil for patch disease					
Current height	0.140"						
Frequency	7 days/week						
Type of mower	walking				OTHER PRODUCTS		
Rollers used	weekly in June - October				Date(s)	Product	Rate (oz./M)
Groomers used							
CULTIVATION		NOTES/COMMENTS				Aquafer Wetting agent	4 oz./K every 30 days
Aerification - dates	5/15 and 8/14	Lots of disease fall of 2000. Tifgreen almost gone - taken over by adjacent plots. High nematode populations on Floradwarf.					
Aerification - type	5/8" Hollow Tines						
Verticutting	Verticut lightly every other						
Dates of topdressing	Monday in June-Oct. & topdressed lightly on the opposite Mondays. Topdressing monthly in November to May.						
Other cultural practices							

Product and company names mentioned herein may be trademarks and/or registered trademarks of their respective owners