

TABLE 2D. 2001 MANAGEMENT - ON-SITE BERMUDAGRASS TEST AT MOBILE, AL (COUNTRY CLUB OF MOBILE)

ESTABLISHMENT		FERTILIZATION			HERBICIDES			
Planting date	18-Jun-98	Date(s)	Product	Rate (lbs./M)	Date(s)	Product	Rate (oz./M)	
Problems during	None							
	1 lb. N/M, amend soil with Ca as needed. Topdress & roll until smooth surface was attained		1 lb. Of N per month 1/2 lb. Of P per month 2 lbs. Of K per month		None			
			Spray minors and wetting agent monthly rates for fertilizer cut by 1/2 in Winter					
FACTORS OF PLAY								
Date opened for play	always open	FUNGICIDES			INSECTICIDES			
Date closed for play		Date(s)	Product	Rate (oz./M)	Date(s)	Product	Rate (oz./M)	
Type of spikes allowed	softspikes							
Uses of green	putting, chipping,		Chipco top choice in June for mole crickets		None			
			Everything else as needed					
MOWING								
Initial height	0.250"							
Current height	0.135"							
Frequency	7 days/week							
Type of mower	Jacobsen 22" walkers				OTHER PRODUCTS			
Rollers used	No				Date(s)	Product	Rate (oz./M)	
Groomers used	No							
CULTIVATION		NOTES/COMMENTS			every 2 weeks in Summer	Primo	2 oz./K	
Aerification - dates	May and August	The green was not overseeded in Winter 2001-20. Some differences in cold tolerance and greenup were noted. Some varieties (Tifgreen and Floradwarf) are almost completely overtaken by surrounding varieties.						
Aerification - type	1/2" Hollow Tines							
Verticutting	Verticut as needed - about 6 times per growing season							
Dates of topdressing	Topdress every Monday lightly. Primo at 2 oz./M every 2 weeks in Summer.							
Other cultural practices	Brushing after topdressing							

Product and company names mentioned herein may be trademarks and/or registered trademarks of their respective owners