

TABLE 4A. 2000-01 MANAGEMENT - ON-SITE OVERSEEDING OF BERMUDAGRASS TEST AT DULUTH (ATLANTA), GA (ATLANTA ATHLETIC CLUB)

SEEDBED PREPARATION

Scalped at 0.25" and handraked. After seeding, brushed with push brooms in two directions.

MOWING

Initial height 0.25"
 mowing height during test 7/16"
 Frequency 3-4 times per week
 Type of mower JD 3235 Reel

CULTIVATION

Aerification - dates 22-Jul-01
 Aerification - type 3/4" solid
 Verticutting
 Dates of topdressing
 Other cultural practices

MAINTENANCE PRACTICES TO ENHANCE SPRING TRANSITION

Solid tine (3/4") on July 22

OTHER INFORMATION

Course closed for overseeding Yes, for 1 day
 Landing area Yes
 Divoting problem No
 Cart allowed on Test area No carts until Bermuda greenup
 Traffic patterns No

FERTILIZATION

Date(s)	Product	Rate (lbs./M)
1 week after seeding	19-19-19	0.25 lb. N
23-Apr-01	19-19-19	1.5 lb. N
5-Jun-01	42-0-0 (multicote)	1 lb. N

FUNGICIDES

Date(s)	Product	Rate (oz./M)
None		

HERBICIDES

Date(s)	Product	Rate (oz./M)
None		

INSECTICIDES

Date(s)	Product	Rate (oz./M)
None		

OTHER PRODUCTS

Date(s)	Product	Rate (oz./M)
None		

NOTES/COMMENTS

Different types of Bermudagrass affected overseeding establishment.

Product and company names mentioned herein may be trademarks and/or registered trademarks of their respective owners